Nurses Content
Downloaded from nursescontent.com

Alcohol withdrawal Initial assessment

Mental Health Nursing | Initial assessment

For educational purposes only.

Fictional educational example. Individualize using verified patient findings, authorized treatment and current local protocols. No real patient record or observed outcome is claimed.

Patient profile and history

43-year-old man admitted after stopping heavy alcohol use. Care stage: At the first supervised assessment.

Medical history: Sustained heavy alcohol use was followed by recent cessation. Tremor, sweating and anxiety developed; previous severe withdrawal is specifically assessed. No known drug allergy is reported in this fictional interview; this does not replace medicine-by-medicine verification in real care.

Surgical history: No previous major operation is reported in this fictional interview. Confirm procedure details, complications and any relevant devices when applying the exercise.

Personal and family assessment: The person wishes to participate in decisions and prefers permission before others are involved. Assess support, work/study function, substance use and practical access without stigma.

Scenario: The patient or caregiver wants to know which concern needs attention first.

Clinical assessment

Subjective: Tremor, sweating, anxiety and poor sleep.

Objective findings and required checks: Autonomic symptoms require structured monitoring; seizure and delirium risk are assessed.

Physical examination and investigations: Assess withdrawal severity, observations, mental state, glucose, electrolytes, nutrition and prior severe withdrawal.

Supply actual values, units, timing and trends during supervised application. Do not invent missing clinical data.

Clinical focus and pathophysiology

Alcohol withdrawal: Neuroadaptation to sustained alcohol exposure can produce dangerous hyperexcitability after cessation.

Medical / surgical management

Follow the supervised withdrawal pathway with frequent reassessment, safety support and prescribed treatment.

Nurses Content For educational purposes only. 1 / 7

Nurses Content
Downloaded from nursescontent.com

Illustrative drug and treatment study

Medicine / treatment	Purpose	Nursing checks
Prescribed withdrawal medicine	Control withdrawal under a monitored protocol	Watch sedation, breathing and response
prescribed thiamine	Prevent or treat deficiency-related neurological harm	Follow the route and timing ordered; do not delay emergency glucose for hypoglycemia.

Treatment examples are not a prescription. Verify the authorized product, dose, route, timing, contraindications and current local product information.

Nursing process - ADPIE

Assessment

Tremor, sweating, anxiety and poor sleep. Autonomic symptoms require structured monitoring; seizure and delirium risk are assessed. The patient or caregiver wants to know which concern needs attention first.

Nursing diagnosis

Prioritize vulnerability to injury during seizure activity; immediate safety needs during altered mood, cognition or behaviour; sleep disrupted by symptoms or the care environment; learning needs about the condition and care plan. Validate each concern against assessment. For an actual problem, link the problem, contributing factors and observed cues; for a risk problem, document the risk factors without invented symptoms.

Planning

After the first indicated intervention, at the documented review time: document a patient-specific baseline and measurable target. Agree short- and longer-term outcomes with the patient/caregiver; physiological safety takes priority over teaching and discharge tasks.

Implementation

Follow the supervised withdrawal pathway with frequent reassessment, safety support and prescribed treatment. Establish the immediate clinical priorities and verify the baseline before assuming a diagnosis. Document care, prescribed treatment, consent and escalation.

Evaluation

Compare reassessment with each stated outcome. Mark met, partly met or not met using observed evidence, document the review time and revise the plan. This vignette provides no completed patient outcome.

Nurses Content For educational purposes only. 2 / 7

Nurses Content
Downloaded from nursescontent.com

Nursing care plan

Nursing diagnoses use original clinical problem wording. They are not the licensed NANDA-I classification. Verify any formal label or code against your institution's licensed current edition; select actual or risk diagnoses only when supported by assessment.

Context task: Establish the immediate clinical priorities and verify the baseline before assuming a diagnosis.

Set individualized measurable targets and an exact review time. Select actual or risk statements only when supported by assessment.

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask about warning symptoms, seizure history and prescribed treatment. Assess: Observe consciousness, seizure features, duration and recovery.	Vulnerability to injury during seizure activity. Possible factors: altered neurological activity or the underlying condition. Validate against the patient's assessment.	Short-term: After the first indicated intervention, at the documented review time: The patient is protected from avoidable injury and urgent prolonged activity is escalated. Longer-term: Before the clinically approved transition or agreed follow-up: Patient or caregiver explains the individualized seizure first-aid and follow-up plan.	1. Time and describe seizure activity and call for urgent help according to protocol. Rationale: Duration and recovery help identify an emergency. 2. Protect from nearby hazards, avoid restraint and place nothing in the mouth. Rationale: These actions reduce avoidable trauma and airway obstruction. 3. Support airway assessment, appropriate recovery positioning and prescribed emergency care. Rationale: Breathing and circulation may be compromised during or after a seizure. 4. Follow the supervised withdrawal pathway with frequent reassessment, safety support and prescribed treatment. Rationale: Neuroadaptation to sustained alcohol exposure can produce dangerous hyperexcitability after cessation. The condition-specific plan addresses the underlying problem while response is monitored.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 3 / 7

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask directly about safety, self-harm, aggression and access to harmful means. Assess: Assess risk, observation needs and environmental hazards collaboratively.	Immediate safety needs during altered mood, cognition or behaviour. Possible factors: reduced judgment, severe distress or loss of control. Validate against the patient's assessment.	Short-term: After the first indicated intervention, at the documented review time: Immediate risks are addressed using the least restrictive appropriate care plan. Longer-term: Before the clinically approved transition or agreed follow-up: A collaborative safety and follow-up plan is documented before transition.	1. Escalate imminent danger immediately and maintain observation according to risk and local policy. Rationale: Timely support protects life and prevents unobserved deterioration. 2. Use de-escalation, respectful communication and proportionate environmental safeguards. Rationale: Dignity and reduced stimulation can support safer engagement. 3. Review risk with the multidisciplinary team and document changes and handover. Rationale: Risk is dynamic and requires continuity across staff and settings.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask about usual sleep, current disruption and daytime fatigue. Assess: Record sleep pattern, symptom triggers and environmental disturbance.	Sleep disrupted by symptoms or the care environment. Possible factors: discomfort, anxiety or interrupted routines. Validate against the patient's assessment.	Short-term: After the first indicated intervention, at the documented review time: Reports improved rest during the next planned sleep period. Longer-term: Before the clinically approved transition or agreed follow-up: Identifies practical sleep-supporting routines for recovery.	1. Identify pain, breathlessness, anxiety and medicine-related causes of poor sleep. Rationale: Treating contributors is more useful than assuming a single cause. 2. Cluster non-urgent care and reduce light and noise when safe. Rationale: A predictable environment protects opportunities to sleep. 3. Use relaxation strategies and review prescribed sedating medicines carefully. Rationale: Rest should improve without avoidable oversedation or falls.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask what the learner already knows and wants to	Learning needs about the condition and care plan. Possible	Short-term: After the first indicated intervention, at the	1. Assess language, literacy, hearing, vision and preferred learning method. Rationale: Accessible teaching improves	Record the baseline and reassessed

Nurses Content For educational purposes only. 4 / 7

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
understand. Assess: Use teach-back or demonstration to identify specific gaps.	factors: new information, complex treatment or limited prior teaching. Validate against the patient's assessment.	documented review time: Explains the most important immediate action in their own words. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates the agreed care skills and names follow-up and warning signs.	understanding. 2. Teach a few condition-specific points using plain language and demonstration. Rationale: Focused information is easier to retain and apply. 3. Use teach-back, correct misunderstandings and provide an accessible written plan. Rationale: Checking understanding supports safer self-management. 4. After stabilization discuss relapse-prevention services, nutrition, medicines and follow-up without blame. Rationale: Condition-specific teaching connects the treatment plan to safe continuing care.	finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 5 / 7

Nurses Content
Downloaded from nursescontent.com

Health education and continuing care

After stabilization discuss relapse-prevention services, nutrition, medicines and follow-up without blame.

Transition only after clinical readiness is confirmed. Reconcile medicines, check access and caregiver support, and document who will review the patient and when.

Urgent warning signs

Seizure, delirium, severe agitation, unstable observations or reduced consciousness is an emergency.

Evaluation and conclusion

For each goal, document met, partly met or not met using observed evidence and the review time. Reassess and revise if progress is inadequate. Individual assessment, prioritized care and coordinated review guide this topic.

Quick Revision

- Neuroadaptation to sustained alcohol exposure can produce dangerous hyperexcitability after cessation.

- Priority concerns: Vulnerability to injury during seizure activity; Immediate safety needs during altered mood, cognition or behaviour; Sleep disrupted by symptoms or the care environment; Learning needs about the condition and care plan.

- Seizure, delirium, severe agitation, unstable observations or reduced consciousness is an emergency.

- After stabilization discuss relapse-prevention services, nutrition, medicines and follow-up without blame.

Exam / viva: Which finding changes the order of priorities, and what must be escalated immediately?

References

Nursing Fundamentals, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2024
https://wtcs.pressbooks.pub/nursingfundamentals/

Nursing Mental Health and Community Concepts, 2e - Open RN / WisTech Open, 2025
https://wtcs.pressbooks.pub/nursingmhcc/

Nursing Pharmacology, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2023
https://wtcs.pressbooks.pub/pharmacology2e/

B.Sc. Nursing curriculum and regulatory notifications - Indian Nursing Council
https://www.indiannursingcouncil.org/gazette-notification

Original cases and care-plan wording by Nurses Content. Linked sources support further study; source terms and local clinical requirements apply. Prepared 23 November 2026. Not independently

Nurses Content For educational purposes only. 6 / 7

Nurses Content
Downloaded from nursescontent.com

clinically peer reviewed.

Rights and reuse

Delivered by Nurses Content. Copyright claims apply only to original material owned by its authors. Medical facts, statutory exceptions and third-party/open licence rights remain unaffected. See the website's /policies/reuse/ page for terms.

Nurses Content For educational purposes only. 7 / 7