Nurses Content
Downloaded from nursescontent.com

Childhood iron-deficiency anemia Transition & follow-up

Child Health Nursing | Transition & follow-up

For educational purposes only.

Fictional educational example. Individualize using verified patient findings, authorized treatment and current local protocols. No real patient record or observed outcome is claimed.

Patient profile and history

5-year-old girl with caregiver present. Care stage: During planning for the next care setting after the team confirms clinical readiness.

Medical history: The caregiver reports a limited diet and gradually reduced activity. Blood testing confirms iron deficiency; growth and other causes are reviewed. No known drug allergy is reported in this fictional interview; this does not replace medicine-by-medicine verification in real care.

Surgical history: No previous major operation is reported in this fictional interview. Confirm procedure details, complications and any relevant devices when applying the exercise.

Personal and family assessment: A caregiver is available for teaching and assisted care. The family requests plain-language instructions, a practical feeding/activity plan and follow-up. Assess development, immunization record and safeguarding individually.

Scenario: The next care setting needs a clear handover and the patient or caregiver wants to know when to seek urgent review.

Clinical assessment

Subjective: Fatigue, picky eating and reduced playfulness.

Objective findings and required checks: Pallor and laboratory-confirmed iron deficiency; growth and diet require review.

Physical examination and investigations: Review blood count, iron assessment, growth, diet and evaluation for blood loss or other causes.

Supply actual values, units, timing and trends during supervised application. Do not invent missing clinical data.

Clinical focus and pathophysiology

Childhood iron-deficiency anemia: Insufficient iron limits hemoglobin production and oxygen delivery.

Nurses Content For educational purposes only. 1 / 7

Nurses Content
Downloaded from nursescontent.com

Medical / surgical management

Support prescribed iron replacement, suitable diet and follow-up to confirm response.

Illustrative drug and treatment study

Medicine / treatment	Purpose	Nursing checks
Prescribed pediatric iron	Replace iron stores	Verify weight-based prescription, administration advice and safe storage
no unsupervised dose changes	Prevent poisoning or undertreatment	Keep the product out of children's reach.

Treatment examples are not a prescription. Verify the authorized product, dose, route, timing, contraindications and current local product information.

Nursing process - ADPIE

Assessment

Fatigue, picky eating and reduced playfulness. Pallor and laboratory-confirmed iron deficiency; growth and diet require review. The next care setting needs a clear handover and the patient or caregiver wants to know when to seek urgent review.

Nursing diagnosis

Prioritize nutritional intake below current needs; movement limited by weakness, discomfort or restriction; reduced circulation to vulnerable tissues; learning needs about the condition and care plan. Validate each concern against assessment. For an actual problem, link the problem, contributing factors and observed cues; for a risk problem, document the risk factors without invented symptoms.

Planning

Before any clinically approved transition: document a patient-specific baseline and measurable target. Agree short- and longer-term outcomes with the patient/caregiver; physiological safety takes priority over teaching and discharge tasks.

Implementation

Support prescribed iron replacement, suitable diet and follow-up to confirm response. Check unresolved risks, medicine reconciliation, practical support and follow-up before transition. Document care, prescribed treatment, consent and escalation.

Evaluation

Compare reassessment with each stated outcome. Mark met, partly met or not met using observed evidence, document the review time and revise the plan. This vignette provides no completed patient outcome.

Nurses Content For educational purposes only. 2 / 7

Nurses Content
Downloaded from nursescontent.com

Nursing care plan

Nursing diagnoses use original clinical problem wording. They are not the licensed NANDA-I classification. Verify any formal label or code against your institution's licensed current edition; select actual or risk diagnoses only when supported by assessment.

Context task: Check unresolved risks, medicine reconciliation, practical support and follow-up before transition.

Set individualized measurable targets and an exact review time. Select actual or risk statements only when supported by assessment.

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask about usual diet, recent intake, weight change and food access. Assess: Assess growth or weight trends, intake record and relevant nutritional findings.	Nutritional intake below current needs. Possible factors: poor appetite, increased requirements or feeding difficulty. Validate against the patient's assessment.	Short-term: Before any clinically approved transition: Tolerates the agreed nutrition plan without new complications. Longer-term: Before the clinically approved transition or agreed follow-up: Explains the individualized meal or feeding plan and follow-up arrangements.	1. Assess nutritional risk and confirm swallowing safety and any fasting order. Rationale: The appropriate feeding route depends on clinical safety. 2. Offer suitable small meals or prescribed feeds with dietitian input when indicated. Rationale: Individualized support improves intake and nutrient adequacy. 3. Monitor tolerance, weight or growth and metabolic findings where required. Rationale: Progress and complications guide adjustment of support. 4. Support prescribed iron replacement, suitable diet and follow-up to confirm response. Rationale: Insufficient iron limits hemoglobin production and oxygen delivery. The condition-specific plan addresses the underlying problem while response is monitored.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask about usual mobility, fatigue and help needed with	Movement limited by weakness, discomfort or restriction.	Short-term: Before any clinically approved transition: Completes a	1. Assess assistance needs and confirm prescribed restrictions before mobilization. Rationale: Matching	Record the baseline and reassessed

Nurses Content For educational purposes only. 3 / 7

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
activities. Assess: Assess strength, range, gait, balance and ordered movement restrictions.	Possible factors: reduced strength, pain or prescribed movement limits. Validate against the patient's assessment.	jointly agreed movement task safely with appropriate assistance. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates the home exercise or mobility plan before discharge.	support to ability prevents injury. 2. Plan graded movement, position changes and rest with physiotherapy where indicated. Rationale: Progressive activity supports function while limiting exhaustion. 3. Monitor tolerance and stop for new concerning symptoms. Rationale: Symptoms may indicate unsafe exertion or deterioration.	finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask about dizziness, chest discomfort, fatigue and cold extremities. Assess: Assess consciousness, pulse, blood pressure, capillary refill and urine trend.	Reduced circulation to vulnerable tissues. Possible factors: altered blood flow or cardiac function. Validate against the patient's assessment.	Short-term: Before any clinically approved transition: Circulation and consciousness remain within the clinician-agreed targets. Longer-term: Before the clinically approved transition or agreed follow-up: Describes symptoms that require immediate review and the agreed follow-up plan.	1. Trend perfusion findings and escalate new chest pain, shock or altered consciousness immediately. Rationale: Poor circulation can rapidly threaten organ function. 2. Balance activity and rest and maintain the ordered monitoring schedule. Rationale: Reducing demand supports recovery while trends guide treatment. 3. Administer prescribed therapy and monitor effects and adverse reactions. Rationale: Response monitoring supports safe correction of the underlying problem.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask what the learner already knows and wants to understand. Assess: Use teach-back or demonstration to identify specific	Learning needs about the condition and care plan. Possible factors: new information, complex treatment or limited prior teaching.	Short-term: Before any clinically approved transition: Explains the most important immediate action in their own words. Longer-term: Before the clinically	1. Assess language, literacy, hearing, vision and preferred learning method. Rationale: Accessible teaching improves understanding. 2. Teach a few condition-specific points using plain language and demonstration. Rationale: Focused information is easier to retain	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not

Nurses Content For educational purposes only. 4 / 7

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
gaps.	Validate against the patient's assessment.	approved transition or agreed follow-up: Demonstrates the agreed care skills and names follow-up and warning signs.	and apply. 3. Use teach-back, correct misunderstandings and provide an accessible written plan. Rationale: Checking understanding supports safer self-management. 4. Teach iron-containing foods, correct medicine measurement, expected effects and follow-up testing. Rationale: Condition-specific teaching connects the treatment plan to safe continuing care.	met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 5 / 7

Nurses Content
Downloaded from nursescontent.com

Health education and continuing care

Teach iron-containing foods, correct medicine measurement, expected effects and follow-up testing.

Transition only after clinical readiness is confirmed. Reconcile medicines, check access and caregiver support, and document who will review the patient and when.

Urgent warning signs

Marked breathlessness, syncope, cardiac symptoms or suspected accidental iron ingestion requires urgent care.

Evaluation and conclusion

For each goal, document met, partly met or not met using observed evidence and the review time. Reassess and revise if progress is inadequate. Individual assessment, prioritized care and coordinated review guide this topic.

Quick Revision

- Insufficient iron limits hemoglobin production and oxygen delivery.

- Priority concerns: Nutritional intake below current needs; Movement limited by weakness, discomfort or restriction; Reduced circulation to vulnerable tissues; Learning needs about the condition and care plan.

- Marked breathlessness, syncope, cardiac symptoms or suspected accidental iron ingestion requires urgent care.

- Teach iron-containing foods, correct medicine measurement, expected effects and follow-up testing.

Exam / viva: Which unresolved problem would delay transition, and what information must reach the next care team?

References

Nursing Fundamentals, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2024
https://wtcs.pressbooks.pub/nursingfundamentals/

Health conditions A to Z: external further reading - NHS
https://www.nhs.uk/conditions/

Nursing Pharmacology, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2023
https://wtcs.pressbooks.pub/pharmacology2e/

B.Sc. Nursing curriculum and regulatory notifications - Indian Nursing Council
https://www.indiannursingcouncil.org/gazette-notification

Original cases and care-plan wording by Nurses Content. Linked sources support further study; source terms and local clinical requirements apply. Prepared 23 November 2026. Not independently clinically peer reviewed.

Nurses Content For educational purposes only. 6 / 7

Nurses Content
Downloaded from nursescontent.com

Rights and reuse

Delivered by Nurses Content. Copyright claims apply only to original material owned by its authors. Medical facts, statutory exceptions and third-party/open licence rights remain unaffected. See the website's /policies/reuse/ page for terms.

Nurses Content For educational purposes only. 7 / 7