Nurses Content
Downloaded from nursescontent.com

Palliative care at home Transition & follow-up

Community Health Nursing | Transition & follow-up

For educational purposes only.

Fictional educational example. Individualize using verified patient findings, authorized treatment and current local protocols. No real patient record or observed outcome is claimed.

Patient profile and history

61-year-old man with advanced illness and a home-care plan. Care stage: During planning for the next care setting after the team confirms clinical readiness.

Medical history: Advanced illness is followed by a palliative-care team. Comfort, dignity and the person's documented goals guide the home-care plan. No known drug allergy is reported in this fictional interview; this does not replace medicine-by-medicine verification in real care.

Surgical history: No previous major operation is reported in this fictional interview. Confirm procedure details, complications and any relevant devices when applying the exercise.

Personal and family assessment: The patient wants to regain the usual level of independence. Review available support, daily activities, treatment access and preferred language rather than assuming these are the same for every person.

Scenario: The next care setting needs a clear handover and the patient or caregiver wants to know when to seek urgent review.

Clinical assessment

Subjective: Pain, fatigue and concern about burdening family.

Objective findings and required checks: Symptom and caregiver needs require review; goals of care guide interventions.

Physical examination and investigations: Assess symptoms, function, medicines, caregiver capacity and the documented goals and escalation plan.

Supply actual values, units, timing and trends during supervised application. Do not invent missing clinical data.

Clinical focus and pathophysiology

Palliative care at home: Advanced illness can create physical symptoms and emotional, social and spiritual needs.

Nurses Content For educational purposes only. 1 / 8

Nurses Content
Downloaded from nursescontent.com

Medical / surgical management

Prioritize comfort, dignity, shared decisions and coordinated home/palliative-team support.

Illustrative drug and treatment study

Medicine / treatment	Purpose	Nursing checks
Prescribed analgesic	Relieve pain according to the individualized palliative plan	Monitor effect, sedation, breathing and bowel needs
prescribed symptom medicine	Treat a defined symptom	Confirm route and caregiver instructions.

Treatment examples are not a prescription. Verify the authorized product, dose, route, timing, contraindications and current local product information.

Nursing process - ADPIE

Assessment

Pain, fatigue and concern about burdening family. Symptom and caregiver needs require review; goals of care guide interventions. The next care setting needs a clear handover and the patient or caregiver wants to know when to seek urgent review.

Nursing diagnosis

Prioritize pain interfering with rest or activity; daily self-care needs exceed current ability; coping resources strained by illness or social stress; learning needs about the condition and care plan. Validate each concern against assessment. For an actual problem, link the problem, contributing factors and observed cues; for a risk problem, document the risk factors without invented symptoms.

Planning

Before any clinically approved transition: document a patient-specific baseline and measurable target. Agree short- and longer-term outcomes with the patient/caregiver; physiological safety takes priority over teaching and discharge tasks.

Implementation

Prioritize comfort, dignity, shared decisions and coordinated home/palliative-team support. Check unresolved risks, medicine reconciliation, practical support and follow-up before transition. Document care, prescribed treatment, consent and escalation.

Evaluation

Compare reassessment with each stated outcome. Mark met, partly met or not met using observed evidence, document the review time and revise the plan. This

Nurses Content For educational purposes only. 2 / 8

Nurses Content
Downloaded from nursescontent.com

vignette provides no completed patient outcome.

Nurses Content For educational purposes only. 3 / 8

Nurses Content
Downloaded from nursescontent.com

Nursing care plan

Nursing diagnoses use original clinical problem wording. They are not the licensed NANDA-I classification. Verify any formal label or code against your institution's licensed current edition; select actual or risk diagnoses only when supported by assessment.

Context task: Check unresolved risks, medicine reconciliation, practical support and follow-up before transition.

Set individualized measurable targets and an exact review time. Select actual or risk statements only when supported by assessment.

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask about pain site, onset, severity, quality and the patient's acceptable pain level. Assess: Observe guarding, expression and effect on movement; use an appropriate pain scale.	Pain interfering with rest or activity. Possible factors: tissue irritation, injury or the underlying condition. Validate against the patient's assessment.	Short-term: Before any clinically approved transition: Pain decreases toward the patient-agreed target after reassessment. Longer-term: Before the clinically approved transition or agreed follow-up: Uses agreed comfort strategies and explains when new or worsening pain needs review.	1. Assess pain and associated red flags before and after treatment. Rationale: Changing pain may signal a complication and helps evaluate relief. 2. Provide comfortable positioning, reassurance and appropriate non-drug measures. Rationale: Comfort measures can reduce distress and complement prescribed treatment. 3. Give prescribed analgesia and monitor sedation, breathing and other adverse effects. Rationale: Relief should improve function without creating medication-related harm. 4. Prioritize comfort, dignity, shared decisions and coordinated home/palliative-team support. Rationale: Advanced illness can create physical symptoms and emotional, social and spiritual needs. The condition-specific plan addresses the underlying problem while response is monitored.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 4 / 8

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask which activities the person can do and where help is wanted. Assess: Assess personal-care tasks, cognition, mobility and available support.	Daily self-care needs exceed current ability. Possible factors: weakness, symptoms or reduced confidence. Validate against the patient's assessment.	Short-term: Before any clinically approved transition: Completes a chosen self-care activity with the agreed level of assistance. Longer-term: Before the clinically approved transition or agreed follow-up: Patient or caregiver demonstrates essential care skills and knows support options.	1. Assess ability task by task and respect preferences and privacy. Rationale: Specific assessment prevents unnecessary dependence. 2. Provide only the assistance required and use suitable aids. Rationale: Supported participation protects dignity and builds independence. 3. Teach caregivers with consent and confirm realistic support at home. Rationale: Continuity depends on available skills and resources.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask about stressors, strengths, support people and previous coping. Assess: Assess function, engagement and immediate safety needs.	Coping resources strained by illness or social stress. Possible factors: demands exceeding currently available support. Validate against the patient's assessment.	Short-term: Before any clinically approved transition: Chooses one feasible coping action and identifies a support person or service. Longer-term: Before the clinically approved transition or agreed follow-up: Has a realistic follow-up and support plan agreed with the care team.	1. Assess strengths and barriers collaboratively. Rationale: An individualized plan builds on resources the person can actually use. 2. Practice a simple coping strategy and divide tasks into achievable steps. Rationale: Small successes support confidence and participation. 3. Coordinate social, psychological or community support as indicated. Rationale: Practical barriers can limit recovery even when clinical treatment is available.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask what the learner already knows and wants to understand. Assess:	Learning needs about the condition and care plan. Possible factors: new	Short-term: Before any clinically approved transition: Explains the most important immediate	1. Assess language, literacy, hearing, vision and preferred learning method. Rationale: Accessible teaching improves understanding. 2. Teach a few	Record the baseline and reassessed finding, the time

Nurses Content For educational purposes only. 5 / 8

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Use teach-back or demonstration to identify specific gaps.	information, complex treatment or limited prior teaching. Validate against the patient's assessment.	action in their own words. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates the agreed care skills and names follow-up and warning signs.	condition-specific points using plain language and demonstration. Rationale: Focused information is easier to retain and apply. 3. Use teach-back, correct misunderstandings and provide an accessible written plan. Rationale: Checking understanding supports safer self-management. 4. Teach medicine administration only as authorized, comfort measures, support contacts and anticipatory planning. Rationale: Condition-specific teaching connects the treatment plan to safe continuing care.	and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 6 / 8

Nurses Content
Downloaded from nursescontent.com

Health education and continuing care

Teach medicine administration only as authorized, comfort measures, support contacts and anticipatory planning.

Transition only after clinical readiness is confirmed. Reconcile medicines, check access and caregiver support, and document who will review the patient and when.

Urgent warning signs

Uncontrolled symptoms, unexpected deterioration or caregiver inability to provide safe care requires review according to the agreed escalation plan.

Evaluation and conclusion

For each goal, document met, partly met or not met using observed evidence and the review time. Reassess and revise if progress is inadequate. Individual assessment, prioritized care and coordinated review guide this topic.

Quick Revision

- Advanced illness can create physical symptoms and emotional, social and spiritual needs.

- Priority concerns: Pain interfering with rest or activity; Daily self-care needs exceed current ability; Coping resources strained by illness or social stress; Learning needs about the condition and care plan.

- Uncontrolled symptoms, unexpected deterioration or caregiver inability to provide safe care requires review according to the agreed escalation plan.

- Teach medicine administration only as authorized, comfort measures, support contacts and anticipatory planning.

Exam / viva: Which unresolved problem would delay transition, and what information must reach the next care team?

References

Nursing Fundamentals, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2024
https://wtcs.pressbooks.pub/nursingfundamentals/

Nursing Mental Health and Community Concepts, 2e - Open RN / WisTech Open, 2025
https://wtcs.pressbooks.pub/nursingmhcc/

Nursing Pharmacology, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2023
https://wtcs.pressbooks.pub/pharmacology2e/

B.Sc. Nursing curriculum and regulatory notifications - Indian Nursing Council
https://www.indiannursingcouncil.org/gazette-notification

Original cases and care-plan wording by Nurses Content. Linked sources support further study; source terms and local clinical requirements apply. Prepared 23 November 2026. Not independently clinically peer reviewed.

Nurses Content For educational purposes only. 7 / 8

Nurses Content
Downloaded from nursescontent.com

Rights and reuse

Delivered by Nurses Content. Copyright claims apply only to original material owned by its authors. Medical facts, statutory exceptions and third-party/open licence rights remain unaffected. See the website's /policies/reuse/ page for terms.

Nurses Content For educational purposes only. 8 / 8