Nurses Content
Downloaded from nursescontent.com

Perioperative anxiety Response monitoring

Nursing Foundation | Response monitoring

For educational purposes only.

Fictional educational example. Individualize using verified patient findings, authorized treatment and current local protocols. No real patient record or observed outcome is claimed.

Patient profile and history

29-year-old woman awaiting planned surgery. Care stage: During review after prescribed initial care.

Medical history: The patient is awaiting the planned operation described in the care episode. Uncertainty about anesthesia and recovery is the main reported source of distress. No known drug allergy is reported in this fictional interview; this does not replace medicine-by-medicine verification in real care.

Surgical history: No previous major operation is reported in this fictional interview. Confirm procedure details, complications and any relevant devices when applying the exercise.

Personal and family assessment: The patient wants to regain the usual level of independence. Review available support, daily activities, treatment access and preferred language rather than assuming these are the same for every person.

Scenario: The team requests a clear account of changes after the prescribed care, rather than a statement that the patient is simply better.

Clinical assessment

Subjective: Worried about anesthesia, pain and being away from family.

Objective findings and required checks: Alert and able to consent; anxious expression and repeated questions.

Physical examination and investigations: Assess concerns, understanding, consent status, allergies and preoperative checklist completion.

Supply actual values, units, timing and trends during supervised application. Do not invent missing clinical data.

Clinical focus and pathophysiology

Perioperative anxiety: Uncertainty and perceived threat can increase distress and reduce information retention.

Nurses Content For educational purposes only. 1 / 8

Nurses Content
Downloaded from nursescontent.com

Medical / surgical management

Provide clear information, privacy, reassurance grounded in facts and referral for unanswered consent questions.

Illustrative drug and treatment study

Medicine / treatment	Purpose	Nursing checks
Prescribed premedication	Use only when ordered as part of the anesthesia plan	Check consent timing, monitoring and sedation risk
no self-medication	Avoid unapproved medicines before surgery	Reconcile all products with the team.

Treatment examples are not a prescription. Verify the authorized product, dose, route, timing, contraindications and current local product information.

Nursing process - ADPIE

Assessment

Worried about anesthesia, pain and being away from family. Alert and able to consent; anxious expression and repeated questions. The team requests a clear account of changes after the prescribed care, rather than a statement that the patient is simply better.

Nursing diagnosis

Prioritize emotional distress affecting participation in care; learning needs about the condition and care plan; sleep disrupted by symptoms or the care environment; immediate safety needs during altered mood, cognition or behaviour. Validate each concern against assessment. For an actual problem, link the problem, contributing factors and observed cues; for a risk problem, document the risk factors without invented symptoms.

Planning

At the next scheduled reassessment, with its interval chosen for acuity: document a patient-specific baseline and measurable target. Agree short- and longer-term outcomes with the patient/caregiver; physiological safety takes priority over teaching and discharge tasks.

Implementation

Provide clear information, privacy, reassurance grounded in facts and referral for unanswered consent questions. Compare the current assessment with the documented baseline and identify benefit, harm or lack of response. Document care, prescribed treatment, consent and escalation.

Evaluation

Compare reassessment with each stated outcome. Mark met, partly met or not met

Nurses Content For educational purposes only. 2 / 8

Nurses Content
Downloaded from nursescontent.com

using observed evidence, document the review time and revise the plan. This vignette provides no completed patient outcome.

Nurses Content For educational purposes only. 3 / 8

Nurses Content
Downloaded from nursescontent.com

Nursing care plan

Nursing diagnoses use original clinical problem wording. They are not the licensed NANDA-I classification. Verify any formal label or code against your institution's licensed current edition; select actual or risk diagnoses only when supported by assessment.

Context task: Compare the current assessment with the documented baseline and identify benefit, harm or lack of response.

Set individualized measurable targets and an exact review time. Select actual or risk statements only when supported by assessment.

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask what worries the patient most and how they prefer to receive information. Assess: Observe distress, attention and ability to engage without making assumptions.	Emotional distress affecting participation in care. Possible factors: uncertainty, symptoms or unfamiliar treatment. Validate against the patient's assessment.	Short-term: At the next scheduled reassessment, with its interval chosen for acuity: Identifies one immediate concern and participates in an agreed next step. Longer-term: Before the clinically approved transition or agreed follow-up: Describes helpful support and coping strategies before transition of care.	1. Listen without judgment and acknowledge specific concerns. Rationale: Feeling heard supports trust and therapeutic engagement. 2. Explain one step at a time and use teach-back with permission. Rationale: Manageable information reduces uncertainty and checks understanding. 3. Offer culturally appropriate support and refer when distress persists or safety concerns arise. Rationale: Additional support may be needed beyond bedside reassurance. 4. Provide clear information, privacy, reassurance grounded in facts and referral for unanswered consent questions. Rationale: Uncertainty and perceived threat can increase distress and reduce information retention. The condition-specific plan addresses the underlying problem while response is monitored.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 4 / 8

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask what the learner already knows and wants to understand. Assess: Use teach-back or demonstration to identify specific gaps.	Learning needs about the condition and care plan. Possible factors: new information, complex treatment or limited prior teaching. Validate against the patient's assessment.	Short-term: At the next scheduled reassessment, with its interval chosen for acuity: Explains the most important immediate action in their own words. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates the agreed care skills and names follow-up and warning signs.	1. Assess language, literacy, hearing, vision and preferred learning method. Rationale: Accessible teaching improves understanding. 2. Teach a few condition-specific points using plain language and demonstration. Rationale: Focused information is easier to retain and apply. 3. Use teach-back, correct misunderstandings and provide an accessible written plan. Rationale: Checking understanding supports safer self-management. 4. Explain preparation, fasting according to the anesthesia instructions, postoperative care and how to ask for help. Rationale: Condition-specific teaching connects the treatment plan to safe continuing care.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask about usual sleep, current disruption and daytime fatigue. Assess: Record sleep pattern, symptom triggers and environmental disturbance.	Sleep disrupted by symptoms or the care environment. Possible factors: discomfort, anxiety or interrupted routines. Validate against the patient's assessment.	Short-term: At the next scheduled reassessment, with its interval chosen for acuity: Reports improved rest during the next planned sleep period. Longer-term: Before the clinically approved transition or agreed follow-up: Identifies practical sleep-supporting routines for recovery.	1. Identify pain, breathlessness, anxiety and medicine-related causes of poor sleep. Rationale: Treating contributors is more useful than assuming a single cause. 2. Cluster non-urgent care and reduce light and noise when safe. Rationale: A predictable environment protects opportunities to sleep. 3. Use relaxation strategies and review prescribed sedating medicines carefully. Rationale: Rest should improve without avoidable oversedation or falls.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 5 / 8

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask directly about safety, self-harm, aggression and access to harmful means. Assess: Assess risk, observation needs and environmental hazards collaboratively.	Immediate safety needs during altered mood, cognition or behaviour. Possible factors: reduced judgment, severe distress or loss of control. Validate against the patient's assessment.	Short-term: At the next scheduled reassessment, with its interval chosen for acuity: Immediate risks are addressed using the least restrictive appropriate care plan. Longer-term: Before the clinically approved transition or agreed follow-up: A collaborative safety and follow-up plan is documented before transition.	1. Escalate imminent danger immediately and maintain observation according to risk and local policy. Rationale: Timely support protects life and prevents unobserved deterioration. 2. Use de-escalation, respectful communication and proportionate environmental safeguards. Rationale: Dignity and reduced stimulation can support safer engagement. 3. Review risk with the multidisciplinary team and document changes and handover. Rationale: Risk is dynamic and requires continuity across staff and settings.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 6 / 8

Nurses Content
Downloaded from nursescontent.com

Health education and continuing care

Explain preparation, fasting according to the anesthesia instructions, postoperative care and how to ask for help.

Transition only after clinical readiness is confirmed. Reconcile medicines, check access and caregiver support, and document who will review the patient and when.

Urgent warning signs

New chest pain, breathing difficulty, allergy symptoms or withdrawal of consent requires immediate communication.

Evaluation and conclusion

For each goal, document met, partly met or not met using observed evidence and the review time. Reassess and revise if progress is inadequate. Individual assessment, prioritized care and coordinated review guide this topic.

Quick Revision

- Uncertainty and perceived threat can increase distress and reduce information retention.

- Priority concerns: Emotional distress affecting participation in care; Learning needs about the condition and care plan; Sleep disrupted by symptoms or the care environment; Immediate safety needs during altered mood, cognition or behaviour.

- New chest pain, breathing difficulty, allergy symptoms or withdrawal of consent requires immediate communication.

- Explain preparation, fasting according to the anesthesia instructions, postoperative care and how to ask for help.

Exam / viva: Which observations demonstrate response, and which findings would require revising or stopping the current intervention?

References

Nursing Fundamentals, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2024
https://wtcs.pressbooks.pub/nursingfundamentals/

Nursing Pharmacology, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2023
https://wtcs.pressbooks.pub/pharmacology2e/

B.Sc. Nursing curriculum and regulatory notifications - Indian Nursing Council
https://www.indiannursingcouncil.org/gazette-notification

Original cases and care-plan wording by Nurses Content. Linked sources support further study; source terms and local clinical requirements apply. Prepared 23 November 2026. Not independently clinically peer reviewed.

Nurses Content For educational purposes only. 7 / 8

Nurses Content
Downloaded from nursescontent.com

Rights and reuse

Delivered by Nurses Content. Copyright claims apply only to original material owned by its authors. Medical facts, statutory exceptions and third-party/open licence rights remain unaffected. See the website's /policies/reuse/ page for terms.

Nurses Content For educational purposes only. 8 / 8