Nurses Content
Downloaded from nursescontent.com

Postpartum uterine infection Comfort & daily function

Midwifery & O.G. Nursing | Comfort & daily function

For educational purposes only.

Fictional educational example. Individualize using verified patient findings, authorized treatment and current local protocols. No real patient record or observed outcome is claimed.

Patient profile and history

29-year-old woman with postpartum fever. Care stage: During supported recovery and essential daily activity.

Medical history: Fictional obstetric history: G2P2 during the first postpartum week. Fever, uterine pain and abnormal lochia developed after the current birth. No known drug allergy is reported in this fictional interview; this does not replace medicine-by-medicine verification in real care.

Surgical history: No previous major operation is reported in this fictional interview. Confirm procedure details, complications and any relevant devices when applying the exercise.

Personal and family assessment: The woman requests respectful explanation and involvement of her chosen support person with consent. Assess access to maternity follow-up, nutrition, transport and infant-care support.

Scenario: The patient or caregiver asks how to manage essential daily care without provoking symptoms or unsafe fatigue.

Clinical assessment

Subjective: Lower abdominal pain, feeling unwell and abnormal-smelling lochia.

Objective findings and required checks: Fever and uterine tenderness need urgent maternity assessment and sepsis screening.

Physical examination and investigations: Review observations, blood tests, cultures when indicated and assessment for retained tissue or other sources.

Supply actual values, units, timing and trends during supervised application. Do not invent missing clinical data.

Clinical focus and pathophysiology

Postpartum uterine infection: Ascending or procedure-associated infection can inflame postpartum uterine tissues and progress systemically.

Nurses Content For educational purposes only. 1 / 7

Nurses Content
Downloaded from nursescontent.com

Medical / surgical management

Escalate promptly, monitor perfusion and provide prescribed antimicrobial and source-directed care.

Illustrative drug and treatment study

Medicine / treatment	Purpose	Nursing checks
Prescribed broad-spectrum antimicrobial	Treat the assessed postpartum infection	Check allergy, timing and response
prescribed analgesic	Relieve discomfort	Reassess pain and clinical progression.

Treatment examples are not a prescription. Verify the authorized product, dose, route, timing, contraindications and current local product information.

Nursing process - ADPIE

Assessment

Lower abdominal pain, feeling unwell and abnormal-smelling lochia. Fever and uterine tenderness need urgent maternity assessment and sepsis screening. The patient or caregiver asks how to manage essential daily care without provoking symptoms or unsafe fatigue.

Nursing diagnosis

Prioritize vulnerability to infection or worsening infection; raised temperature increasing discomfort and fluid needs; pain interfering with rest or activity; learning needs about the condition and care plan. Validate each concern against assessment. For an actual problem, link the problem, contributing factors and observed cues; for a risk problem, document the risk factors without invented symptoms.

Planning

During the next agreed care or activity session: document a patient-specific baseline and measurable target. Agree short- and longer-term outcomes with the patient/caregiver; physiological safety takes priority over teaching and discharge tasks.

Implementation

Escalate promptly, monitor perfusion and provide prescribed antimicrobial and source-directed care. Identify the safest achievable activity or comfort goal without overlooking the underlying illness. Document care, prescribed treatment, consent and escalation.

Evaluation

Compare reassessment with each stated outcome. Mark met, partly met or not met using observed evidence, document the review time and revise the plan. This vignette provides no completed patient outcome.

Nurses Content For educational purposes only. 2 / 7

Nurses Content
Downloaded from nursescontent.com

Nursing care plan

Nursing diagnoses use original clinical problem wording. They are not the licensed NANDA-I classification. Verify any formal label or code against your institution's licensed current edition; select actual or risk diagnoses only when supported by assessment.

Context task: Identify the safest achievable activity or comfort goal without overlooking the underlying illness.

Set individualized measurable targets and an exact review time. Select actual or risk statements only when supported by assessment.

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask about exposure, wound symptoms and recent invasive devices. Assess: Inspect relevant sites and monitor systemic findings and investigation trends.	Vulnerability to infection or worsening infection. Possible factors: disrupted barriers, exposure or reduced host defence. Validate against the patient's assessment.	Short-term: During the next agreed care or activity session: Aseptic care is maintained and any new signs of infection are escalated promptly. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates relevant hygiene and explains the infection-prevention plan.	1. Use hand hygiene and risk-based standard or transmission precautions. Rationale: Interrupting transmission protects the patient, staff and other patients. 2. Review invasive device need with the team and maintain indicated aseptic care. Rationale: Unnecessary devices and barrier breaks increase infection exposure. 3. Monitor for systemic deterioration and administer prescribed antimicrobials on schedule. Rationale: Early escalation and timely indicated treatment reduce avoidable harm. 4. Escalate promptly, monitor perfusion and provide prescribed antimicrobial and source-directed care. Rationale: Ascending or procedure-associated infection can inflame postpartum uterine tissues and progress systemically. The condition-specific plan addresses the underlying problem while response is monitored.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 3 / 7

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask about chills, headache, intake and illness onset. Assess: Record temperature site, trend, hydration and other observations.	Raised temperature increasing discomfort and fluid needs. Possible factors: an inflammatory or infectious process. Validate against the patient's assessment.	Short-term: During the next agreed care or activity session: Temperature-related discomfort improves and clinical deterioration is detected promptly. Longer-term: Before the clinically approved transition or agreed follow-up: Explains monitoring, hydration instructions and warning signs for review.	1. Trend temperature together with circulation, breathing and mental state. Rationale: Temperature alone does not describe illness severity. 2. Provide comfortable clothing and fluids when permitted; avoid extreme cooling. Rationale: Supportive care reduces discomfort without provoking unnecessary stress. 3. Administer prescribed treatment and obtain ordered specimens using correct technique. Rationale: Cause-directed management and reliable samples support recovery.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask about pain site, onset, severity, quality and the patient's acceptable pain level. Assess: Observe guarding, expression and effect on movement; use an appropriate pain scale.	Pain interfering with rest or activity. Possible factors: tissue irritation, injury or the underlying condition. Validate against the patient's assessment.	Short-term: During the next agreed care or activity session: Pain decreases toward the patient-agreed target after reassessment. Longer-term: Before the clinically approved transition or agreed follow-up: Uses agreed comfort strategies and explains when new or worsening pain needs review.	1. Assess pain and associated red flags before and after treatment. Rationale: Changing pain may signal a complication and helps evaluate relief. 2. Provide comfortable positioning, reassurance and appropriate non-drug measures. Rationale: Comfort measures can reduce distress and complement prescribed treatment. 3. Give prescribed analgesia and monitor sedation, breathing and other adverse effects. Rationale: Relief should improve function without creating medication-related harm.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask what the learner already knows and wants to understand. Assess:	Learning needs about the condition and care plan. Possible factors: new	Short-term: During the next agreed care or activity session: Explains the most	1. Assess language, literacy, hearing, vision and preferred learning method. Rationale: Accessible teaching improves understanding. 2. Teach a few	Record the baseline and reassessed finding, the time

Nurses Content For educational purposes only. 4 / 7

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Use teach-back or demonstration to identify specific gaps.	information, complex treatment or limited prior teaching. Validate against the patient's assessment.	important immediate action in their own words. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates the agreed care skills and names follow-up and warning signs.	condition-specific points using plain language and demonstration. Rationale: Focused information is easier to retain and apply. 3. Use teach-back, correct misunderstandings and provide an accessible written plan. Rationale: Checking understanding supports safer self-management. 4. Explain hygiene, prescribed treatment, feeding-related medicine advice and follow-up warning signs. Rationale: Condition-specific teaching connects the treatment plan to safe continuing care.	and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 5 / 7

Nurses Content
Downloaded from nursescontent.com

Health education and continuing care

Explain hygiene, prescribed treatment, feeding-related medicine advice and follow-up warning signs.

Transition only after clinical readiness is confirmed. Reconcile medicines, check access and caregiver support, and document who will review the patient and when.

Urgent warning signs

Shock, confusion, severe abdominal pain, heavy bleeding or respiratory deterioration needs emergency review.

Evaluation and conclusion

For each goal, document met, partly met or not met using observed evidence and the review time. Reassess and revise if progress is inadequate. Individual assessment, prioritized care and coordinated review guide this topic.

Quick Revision

- Ascending or procedure-associated infection can inflame postpartum uterine tissues and progress systemically.

- Priority concerns: Vulnerability to infection or worsening infection; Raised temperature increasing discomfort and fluid needs; Pain interfering with rest or activity; Learning needs about the condition and care plan.

- Shock, confusion, severe abdominal pain, heavy bleeding or respiratory deterioration needs emergency review.

- Explain hygiene, prescribed treatment, feeding-related medicine advice and follow-up warning signs.

Exam / viva: How can care preserve independence while staying within the patient's assessed limits?

References

Nursing Fundamentals, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2024
https://wtcs.pressbooks.pub/nursingfundamentals/

Maternal-Newborn Nursing: external further reading - OpenStax / Rice University
https://openstax.org/books/maternal-newborn-nursing/pages/preface

Nursing Pharmacology, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2023
https://wtcs.pressbooks.pub/pharmacology2e/

B.Sc. Nursing curriculum and regulatory notifications - Indian Nursing Council
https://www.indiannursingcouncil.org/gazette-notification

Original cases and care-plan wording by Nurses Content. Linked sources support further study; source terms and local clinical requirements apply. Prepared 23 November 2026. Not independently clinically peer reviewed.

Nurses Content For educational purposes only. 6 / 7

Nurses Content
Downloaded from nursescontent.com

Rights and reuse

Delivered by Nurses Content. Copyright claims apply only to original material owned by its authors. Medical facts, statutory exceptions and third-party/open licence rights remain unaffected. See the website's /policies/reuse/ page for terms.

Nurses Content For educational purposes only. 7 / 7