Nurses Content
Downloaded from nursescontent.com

Safe insulin administration Initial assessment

Pharmacology-related Cases | Initial assessment

For educational purposes only.

Fictional educational example. Individualize using verified patient findings, authorized treatment and current local protocols. No real patient record or observed outcome is claimed.

Patient profile and history

49-year-old man learning prescribed insulin use. Care stage: At the first supervised assessment.

Medical history: Diabetes requires prescribed insulin. The patient is learning a newly authorized device and needs supervised confirmation of product selection and technique. No known drug allergy is reported in this fictional interview; this does not replace medicine-by-medicine verification in real care.

Surgical history: No previous major operation is reported in this fictional interview. Confirm procedure details, complications and any relevant devices when applying the exercise.

Personal and family assessment: The patient wants to regain the usual level of independence. Review available support, daily activities, treatment access and preferred language rather than assuming these are the same for every person.

Scenario: The patient or caregiver wants to know which concern needs attention first.

Clinical assessment

Subjective: Worried about selecting the right pen and recognizing low glucose.

Objective findings and required checks: Needs supervised return demonstration of preparation, timing and injection technique.

Physical examination and investigations: Review prescription, product/concentration, glucose trend, intake, renal function and injection sites.

Supply actual values, units, timing and trends during supervised application. Do not invent missing clinical data.

Clinical focus and pathophysiology

Safe insulin administration: Insulin lowers glucose; product, dose, timing and intake mismatches can cause serious harm.

Nurses Content For educational purposes only. 1 / 8

Nurses Content
Downloaded from nursescontent.com

Medical / surgical management

Verify the exact insulin order, coordinate with meals as appropriate and use supervised teaching and local checking procedures.

Illustrative drug and treatment study

Medicine / treatment	Purpose	Nursing checks
Prescribed insulin	Replace or supplement insulin action	Verify name, concentration, dose order, route, timing, storage and glucose monitoring
fast-acting glucose	Treat a qualifying low glucose episode safely	Follow the individual rescue plan.

Treatment examples are not a prescription. Verify the authorized product, dose, route, timing, contraindications and current local product information.

Nursing process - ADPIE

Assessment

Worried about selecting the right pen and recognizing low glucose. Needs supervised return demonstration of preparation, timing and injection technique. The patient or caregiver wants to know which concern needs attention first.

Nursing diagnosis

Prioritize blood glucose instability requiring close monitoring; possible treatment-related harm requiring prompt assessment; emotional distress affecting participation in care; learning needs about the condition and care plan. Validate each concern against assessment. For an actual problem, link the problem, contributing factors and observed cues; for a risk problem, document the risk factors without invented symptoms.

Planning

After the first indicated intervention, at the documented review time: document a patient-specific baseline and measurable target. Agree short- and longer-term outcomes with the patient/caregiver; physiological safety takes priority over teaching and discharge tasks.

Implementation

Verify the exact insulin order, coordinate with meals as appropriate and use supervised teaching and local checking procedures. Establish the immediate clinical priorities and verify the baseline before assuming a diagnosis. Document care, prescribed treatment, consent and escalation.

Evaluation

Compare reassessment with each stated outcome. Mark met, partly met or not met using observed evidence, document the review time and revise the plan. This

Nurses Content For educational purposes only. 2 / 8

Nurses Content
Downloaded from nursescontent.com

vignette provides no completed patient outcome.

Nurses Content For educational purposes only. 3 / 8

Nurses Content
Downloaded from nursescontent.com

Nursing care plan

Nursing diagnoses use original clinical problem wording. They are not the licensed NANDA-I classification. Verify any formal label or code against your institution's licensed current edition; select actual or risk diagnoses only when supported by assessment.

Context task: Establish the immediate clinical priorities and verify the baseline before assuming a diagnosis.

Set individualized measurable targets and an exact review time. Select actual or risk statements only when supported by assessment.

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask about intake, usual treatment and symptoms of low or high glucose. Assess: Check glucose using the prescribed schedule and assess consciousness and hydration.	Blood glucose instability requiring close monitoring. Possible factors: altered insulin effect, intake, illness or treatment. Validate against the patient's assessment.	Short-term: After the first indicated intervention, at the documented review time: Glucose moves toward the individualized target without treatment-related hypoglycemia. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates monitoring and explains the personalized sick-day and emergency plan.	1. Verify glucose findings and reassess symptoms rather than treating a number alone. Rationale: Symptoms and trends guide safe interpretation. 2. Coordinate prescribed medication with nutrition and monitor the response. Rationale: Mismatch between treatment and intake can cause dangerous glucose changes. 3. Follow the local emergency pathway for severe hypoglycemia or suspected ketoacidosis. Rationale: These conditions can progress rapidly and require urgent team care. 4. Verify the exact insulin order, coordinate with meals as appropriate and use supervised teaching and local checking procedures. Rationale: Insulin lowers glucose; product, dose, timing and intake mismatches can cause serious harm. The condition-specific plan addresses the underlying problem while response is monitored.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 4 / 8

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask about medicine names, timing, symptoms and non-prescribed products. Assess: Review observations, medication record and relevant drug-specific investigations.	Possible treatment-related harm requiring prompt assessment. Possible factors: adverse medication effects or an administration problem. Validate against the patient's assessment.	Short-term: After the first indicated intervention, at the documented review time: The suspected adverse event is escalated and no further avoidable exposure occurs. Longer-term: Before the clinically approved transition or agreed follow-up: Explains the revised medication plan and warning symptoms before discharge.	1. Assess airway, breathing, circulation and the time course of symptoms. Rationale: Serious adverse reactions require immediate stabilization. 2. Escalate suspected toxicity and follow the authorized hold, monitoring or antidote instructions. Rationale: Treatment decisions depend on the drug and clinical severity. 3. Document and communicate the reaction and reconcile the medicine list. Rationale: Accurate records reduce repeat exposure and support pharmacovigilance.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask what worries the patient most and how they prefer to receive information. Assess: Observe distress, attention and ability to engage without making assumptions.	Emotional distress affecting participation in care. Possible factors: uncertainty, symptoms or unfamiliar treatment. Validate against the patient's assessment.	Short-term: After the first indicated intervention, at the documented review time: Identifies one immediate concern and participates in an agreed next step. Longer-term: Before the clinically approved transition or agreed follow-up: Describes helpful support and coping strategies before transition of care.	1. Listen without judgment and acknowledge specific concerns. Rationale: Feeling heard supports trust and therapeutic engagement. 2. Explain one step at a time and use teach-back with permission. Rationale: Manageable information reduces uncertainty and checks understanding. 3. Offer culturally appropriate support and refer when distress persists or safety concerns arise. Rationale: Additional support may be needed beyond bedside reassurance.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask what the learner already knows and wants to	Learning needs about the condition and care plan. Possible	Short-term: After the first indicated intervention, at the	1. Assess language, literacy, hearing, vision and preferred learning method. Rationale: Accessible teaching improves	Record the baseline and reassessed

Nurses Content For educational purposes only. 5 / 8

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
understand. Assess: Use teach-back or demonstration to identify specific gaps.	factors: new information, complex treatment or limited prior teaching. Validate against the patient's assessment.	documented review time: Explains the most important immediate action in their own words. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates the agreed care skills and names follow-up and warning signs.	understanding. 2. Teach a few condition-specific points using plain language and demonstration. Rationale: Focused information is easier to retain and apply. 3. Use teach-back, correct misunderstandings and provide an accessible written plan. Rationale: Checking understanding supports safer self-management. 4. Teach device technique, site rotation, storage, sharps disposal and the individualized hypoglycemia/sick-day plan. Rationale: Condition-specific teaching connects the treatment plan to safe continuing care.	finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 6 / 8

Nurses Content
Downloaded from nursescontent.com

Health education and continuing care

Teach device technique, site rotation, storage, sharps disposal and the individualized hypoglycemia/sick-day plan.

Transition only after clinical readiness is confirmed. Reconcile medicines, check access and caregiver support, and document who will review the patient and when.

Urgent warning signs

Wrong-product/dose exposure, severe low glucose, altered consciousness or seizure requires immediate help.

Evaluation and conclusion

For each goal, document met, partly met or not met using observed evidence and the review time. Reassess and revise if progress is inadequate. Individual assessment, prioritized care and coordinated review guide this topic.

Quick Revision

- Insulin lowers glucose; product, dose, timing and intake mismatches can cause serious harm.

- Priority concerns: Blood glucose instability requiring close monitoring; Possible treatment-related harm requiring prompt assessment; Emotional distress affecting participation in care; Learning needs about the condition and care plan.

- Wrong-product/dose exposure, severe low glucose, altered consciousness or seizure requires immediate help.

- Teach device technique, site rotation, storage, sharps disposal and the individualized hypoglycemia/sick-day plan.

Exam / viva: Which finding changes the order of priorities, and what must be escalated immediately?

References

Nursing Fundamentals, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2024
https://wtcs.pressbooks.pub/nursingfundamentals/

Nursing Pharmacology, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2023
https://wtcs.pressbooks.pub/pharmacology2e/

B.Sc. Nursing curriculum and regulatory notifications - Indian Nursing Council
https://www.indiannursingcouncil.org/gazette-notification

Original cases and care-plan wording by Nurses Content. Linked sources support further study; source terms and local clinical requirements apply. Prepared 23 November 2026. Not independently clinically peer reviewed.

Nurses Content For educational purposes only. 7 / 8

Nurses Content
Downloaded from nursescontent.com

Rights and reuse

Delivered by Nurses Content. Copyright claims apply only to original material owned by its authors. Medical facts, statutory exceptions and third-party/open licence rights remain unaffected. See the website's /policies/reuse/ page for terms.

Nurses Content For educational purposes only. 8 / 8