Nurses Content
Downloaded from nursescontent.com

Schizophrenia and distressing voices Initial assessment

Mental Health Nursing | Initial assessment

For educational purposes only.

Fictional educational example. Individualize using verified patient findings, authorized treatment and current local protocols. No real patient record or observed outcome is claimed.

Patient profile and history

25-year-old man reporting auditory experiences. Care stage: At the first supervised assessment.

Medical history: The patient has an established psychosis-care history. Distressing voices and reduced self-care have increased; adherence, adverse effects and alternative contributors are reviewed. No known drug allergy is reported in this fictional interview; this does not replace medicine-by-medicine verification in real care.

Surgical history: No previous major operation is reported in this fictional interview. Confirm procedure details, complications and any relevant devices when applying the exercise.

Personal and family assessment: The person wishes to participate in decisions and prefers permission before others are involved. Assess support, work/study function, substance use and practical access without stigma.

Scenario: The patient or caregiver wants to know which concern needs attention first.

Clinical assessment

Subjective: Hears distressing voices and has withdrawn from friends.

Objective findings and required checks: Distractibility and reduced self-care; assess content, distress and immediate risk.

Physical examination and investigations: Assess mental state, risk, physical health, substances and medicine history.

Supply actual values, units, timing and trends during supervised application. Do not invent missing clinical data.

Clinical focus and pathophysiology

Schizophrenia and distressing voices: Psychotic symptoms can alter perception, thought and functioning; causes require comprehensive assessment.

Nurses Content For educational purposes only. 1 / 7

Nurses Content
Downloaded from nursescontent.com

Medical / surgical management

Acknowledge distress without confirming false beliefs, use clear communication and coordinate prescribed psychosocial and medication care.

Illustrative drug and treatment study

Medicine / treatment	Purpose	Nursing checks
Prescribed antipsychotic	Reduce assessed psychotic symptoms	Monitor movement effects, sedation, metabolic parameters and other drug-specific risks
medication review	Identify barriers and adverse effects	Use shared decisions with the treating team.

Treatment examples are not a prescription. Verify the authorized product, dose, route, timing, contraindications and current local product information.

Nursing process - ADPIE

Assessment

Hears distressing voices and has withdrawn from friends. Distractibility and reduced self-care; assess content, distress and immediate risk. The patient or caregiver wants to know which concern needs attention first.

Nursing diagnosis

Prioritize disturbed interpretation of experiences affecting safety or function; immediate safety needs during altered mood, cognition or behaviour; daily self-care needs exceed current ability; learning needs about the condition and care plan. Validate each concern against assessment. For an actual problem, link the problem, contributing factors and observed cues; for a risk problem, document the risk factors without invented symptoms.

Planning

After the first indicated intervention, at the documented review time: document a patient-specific baseline and measurable target. Agree short- and longer-term outcomes with the patient/caregiver; physiological safety takes priority over teaching and discharge tasks.

Implementation

Acknowledge distress without confirming false beliefs, use clear communication and coordinate prescribed psychosocial and medication care. Establish the immediate clinical priorities and verify the baseline before assuming a diagnosis. Document care, prescribed treatment, consent and escalation.

Evaluation

Compare reassessment with each stated outcome. Mark met, partly met or not met using observed evidence, document the review time and revise the plan. This vignette provides no completed patient outcome.

Nurses Content For educational purposes only. 2 / 7

Nurses Content
Downloaded from nursescontent.com

Nursing care plan

Nursing diagnoses use original clinical problem wording. They are not the licensed NANDA-I classification. Verify any formal label or code against your institution's licensed current edition; select actual or risk diagnoses only when supported by assessment.

Context task: Establish the immediate clinical priorities and verify the baseline before assuming a diagnosis.

Set individualized measurable targets and an exact review time. Select actual or risk statements only when supported by assessment.

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask about experiences, distress and thoughts of harming self or others directly and respectfully. Assess: Assess orientation, mental state, behaviour and possible medical contributors.	Disturbed interpretation of experiences affecting safety or function. Possible factors: changes in perception, cognition or mood. Validate against the patient's assessment.	Short-term: After the first indicated intervention, at the documented review time: Engages in a safe, calm interaction and accepts an agreed support step. Longer-term: Before the clinically approved transition or agreed follow-up: Patient and support team describe relapse signs and the follow-up plan.	1. Use calm, brief communication and acknowledge feelings without reinforcing false beliefs. Rationale: A consistent approach reduces confrontation and distress. 2. Assess immediate risk and possible delirium or substance-related causes; escalate concerns. Rationale: Medical and psychiatric emergencies may present with altered behaviour. 3. Maintain a low-stimulation setting and monitor prescribed treatment effects. Rationale: The environment and treatment response influence safety and recovery. 4. Acknowledge distress without confirming false beliefs, use clear communication and coordinate prescribed psychosocial and medication care. Rationale: Psychotic symptoms can alter perception, thought and functioning; causes require comprehensive assessment. The condition-specific plan addresses the underlying problem while response is	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 3 / 7

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
			monitored.	
Ask: Ask directly about safety, self-harm, aggression and access to harmful means. Assess: Assess risk, observation needs and environmental hazards collaboratively.	Immediate safety needs during altered mood, cognition or behaviour. Possible factors: reduced judgment, severe distress or loss of control. Validate against the patient's assessment.	Short-term: After the first indicated intervention, at the documented review time: Immediate risks are addressed using the least restrictive appropriate care plan. Longer-term: Before the clinically approved transition or agreed follow-up: A collaborative safety and follow-up plan is documented before transition.	1. Escalate imminent danger immediately and maintain observation according to risk and local policy. Rationale: Timely support protects life and prevents unobserved deterioration. 2. Use de-escalation, respectful communication and proportionate environmental safeguards. Rationale: Dignity and reduced stimulation can support safer engagement. 3. Review risk with the multidisciplinary team and document changes and handover. Rationale: Risk is dynamic and requires continuity across staff and settings.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask which activities the person can do and where help is wanted. Assess: Assess personal-care tasks, cognition, mobility and available support.	Daily self-care needs exceed current ability. Possible factors: weakness, symptoms or reduced confidence. Validate against the patient's assessment.	Short-term: After the first indicated intervention, at the documented review time: Completes a chosen self-care activity with the agreed level of assistance. Longer-term: Before the clinically approved transition or agreed follow-up: Patient or caregiver demonstrates essential care skills and knows support options.	1. Assess ability task by task and respect preferences and privacy. Rationale: Specific assessment prevents unnecessary dependence. 2. Provide only the assistance required and use suitable aids. Rationale: Supported participation protects dignity and builds independence. 3. Teach caregivers with consent and confirm realistic support at home. Rationale: Continuity depends on available skills and resources.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 4 / 7

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask what the learner already knows and wants to understand. Assess: Use teach-back or demonstration to identify specific gaps.	Learning needs about the condition and care plan. Possible factors: new information, complex treatment or limited prior teaching. Validate against the patient's assessment.	Short-term: After the first indicated intervention, at the documented review time: Explains the most important immediate action in their own words. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates the agreed care skills and names follow-up and warning signs.	1. Assess language, literacy, hearing, vision and preferred learning method. Rationale: Accessible teaching improves understanding. 2. Teach a few condition-specific points using plain language and demonstration. Rationale: Focused information is easier to retain and apply. 3. Use teach-back, correct misunderstandings and provide an accessible written plan. Rationale: Checking understanding supports safer self-management. 4. Teach relapse monitoring, treatment follow-up, physical health care and family support with consent. Rationale: Condition-specific teaching connects the treatment plan to safe continuing care.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 5 / 7

Nurses Content
Downloaded from nursescontent.com

Health education and continuing care

Teach relapse monitoring, treatment follow-up, physical health care and family support with consent.

Transition only after clinical readiness is confirmed. Reconcile medicines, check access and caregiver support, and document who will review the patient and when.

Urgent warning signs

Commands to harm, inability to maintain safety, severe agitation or possible delirium requires urgent assessment.

Evaluation and conclusion

For each goal, document met, partly met or not met using observed evidence and the review time. Reassess and revise if progress is inadequate. Individual assessment, prioritized care and coordinated review guide this topic.

Quick Revision

- Psychotic symptoms can alter perception, thought and functioning; causes require comprehensive assessment.

- Priority concerns: Disturbed interpretation of experiences affecting safety or function; Immediate safety needs during altered mood, cognition or behaviour; Daily self-care needs exceed current ability; Learning needs about the condition and care plan.

- Commands to harm, inability to maintain safety, severe agitation or possible delirium requires urgent assessment.

- Teach relapse monitoring, treatment follow-up, physical health care and family support with consent.

Exam / viva: Which finding changes the order of priorities, and what must be escalated immediately?

References

Nursing Fundamentals, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2024
https://wtcs.pressbooks.pub/nursingfundamentals/

Nursing Mental Health and Community Concepts, 2e - Open RN / WisTech Open, 2025
https://wtcs.pressbooks.pub/nursingmhcc/

Nursing Pharmacology, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2023
https://wtcs.pressbooks.pub/pharmacology2e/

B.Sc. Nursing curriculum and regulatory notifications - Indian Nursing Council
https://www.indiannursingcouncil.org/gazette-notification

Original cases and care-plan wording by Nurses Content. Linked sources support further study; source terms and local clinical requirements apply. Prepared 23 November 2026. Not independently

Nurses Content For educational purposes only. 6 / 7

Nurses Content
Downloaded from nursescontent.com

clinically peer reviewed.

Rights and reuse

Delivered by Nurses Content. Copyright claims apply only to original material owned by its authors. Medical facts, statutory exceptions and third-party/open licence rights remain unaffected. See the website's /policies/reuse/ page for terms.

Nurses Content For educational purposes only. 7 / 7