Nurses Content
Downloaded from nursescontent.com

Type 1 diabetes in childhood Transition & follow-up

Child Health Nursing | Transition & follow-up

For educational purposes only.

Fictional educational example. Individualize using verified patient findings, authorized treatment and current local protocols. No real patient record or observed outcome is claimed.

Patient profile and history

11-year-old girl with caregiver present. Care stage: During planning for the next care setting after the team confirms clinical readiness.

Medical history: Thirst and polyuria led to assessment for type 1 diabetes. The family needs structured insulin, monitoring and school-care education. No known drug allergy is reported in this fictional interview; this does not replace medicine-by-medicine verification in real care.

Surgical history: No previous major operation is reported in this fictional interview. Confirm procedure details, complications and any relevant devices when applying the exercise.

Personal and family assessment: A caregiver is available for teaching and assisted care. The family requests plain-language instructions, a practical feeding/activity plan and follow-up. Assess development, immunization record and safeguarding individually.

Scenario: The next care setting needs a clear handover and the patient or caregiver wants to know when to seek urgent review.

Clinical assessment

Subjective: Thirst, frequent urination and worries about insulin at school.

Objective findings and required checks: Hyperglycemia is under pediatric assessment; ketones and hydration need review.

Physical examination and investigations: Review glucose, ketones when indicated, HbA1c and pediatric diabetes education needs.

Supply actual values, units, timing and trends during supervised application. Do not invent missing clinical data.

Clinical focus and pathophysiology

Type 1 diabetes in childhood: Autoimmune loss of insulin-producing cells creates a need for insulin replacement.

Nurses Content For educational purposes only. 1 / 8

Nurses Content
Downloaded from nursescontent.com

Medical / surgical management

Coordinate the prescribed insulin, meals, activity and monitoring plan with family and school support.

Illustrative drug and treatment study

Medicine / treatment	Purpose	Nursing checks
Prescribed insulin	Replace essential insulin	Verify product, dose order, timing, site rotation and hypoglycemia monitoring
emergency glucose treatment	Treat low glucose according to the child's plan	Confirm safe route and reassessment.

Treatment examples are not a prescription. Verify the authorized product, dose, route, timing, contraindications and current local product information.

Nursing process - ADPIE

Assessment

Thirst, frequent urination and worries about insulin at school. Hyperglycemia is under pediatric assessment; ketones and hydration need review. The next care setting needs a clear handover and the patient or caregiver wants to know when to seek urgent review.

Nursing diagnosis

Prioritize blood glucose instability requiring close monitoring; insufficient fluid replacement for current losses; emotional distress affecting participation in care; learning needs about the condition and care plan. Validate each concern against assessment. For an actual problem, link the problem, contributing factors and observed cues; for a risk problem, document the risk factors without invented symptoms.

Planning

Before any clinically approved transition: document a patient-specific baseline and measurable target. Agree short- and longer-term outcomes with the patient/caregiver; physiological safety takes priority over teaching and discharge tasks.

Implementation

Coordinate the prescribed insulin, meals, activity and monitoring plan with family and school support. Check unresolved risks, medicine reconciliation, practical support and follow-up before transition. Document care, prescribed treatment, consent and escalation.

Evaluation

Compare reassessment with each stated outcome. Mark met, partly met or not met using observed evidence, document the review time and revise the plan. This

Nurses Content For educational purposes only. 2 / 8

Nurses Content
Downloaded from nursescontent.com

vignette provides no completed patient outcome.

Nurses Content For educational purposes only. 3 / 8

Nurses Content
Downloaded from nursescontent.com

Nursing care plan

Nursing diagnoses use original clinical problem wording. They are not the licensed NANDA-I classification. Verify any formal label or code against your institution's licensed current edition; select actual or risk diagnoses only when supported by assessment.

Context task: Check unresolved risks, medicine reconciliation, practical support and follow-up before transition.

Set individualized measurable targets and an exact review time. Select actual or risk statements only when supported by assessment.

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask about intake, usual treatment and symptoms of low or high glucose. Assess: Check glucose using the prescribed schedule and assess consciousness and hydration.	Blood glucose instability requiring close monitoring. Possible factors: altered insulin effect, intake, illness or treatment. Validate against the patient's assessment.	Short-term: Before any clinically approved transition: Glucose moves toward the individualized target without treatment-related hypoglycemia. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates monitoring and explains the personalized sick-day and emergency plan.	1. Verify glucose findings and reassess symptoms rather than treating a number alone. Rationale: Symptoms and trends guide safe interpretation. 2. Coordinate prescribed medication with nutrition and monitor the response. Rationale: Mismatch between treatment and intake can cause dangerous glucose changes. 3. Follow the local emergency pathway for severe hypoglycemia or suspected ketoacidosis. Rationale: These conditions can progress rapidly and require urgent team care. 4. Coordinate the prescribed insulin, meals, activity and monitoring plan with family and school support. Rationale: Autoimmune loss of insulin-producing cells creates a need for insulin replacement. The condition-specific plan addresses the underlying problem while response is monitored.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 4 / 8

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask about thirst, intake, vomiting, stools and dizziness. Assess: Monitor weight trend, intake/output, mucosa, circulation and relevant electrolytes.	Insufficient fluid replacement for current losses. Possible factors: intake below requirements or increased fluid loss. Validate against the patient's assessment.	Short-term: Before any clinically approved transition: Hydration and circulation improve within the agreed reassessment interval. Longer-term: Before the clinically approved transition or agreed follow-up: Patient or caregiver explains the individualized fluid plan and warning signs.	1. Measure intake and losses using age-appropriate methods. Rationale: Accurate trends guide replacement and help prevent overcorrection. 2. Offer oral fluids only when swallowing, consciousness and the treatment plan permit; otherwise implement prescribed replacement. Rationale: The safest route depends on aspiration risk and clinical severity. 3. Reassess perfusion, weight and laboratory trends after prescribed replacement. Rationale: Both continued dehydration and fluid overload require prompt recognition.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask what worries the patient most and how they prefer to receive information. Assess: Observe distress, attention and ability to engage without making assumptions.	Emotional distress affecting participation in care. Possible factors: uncertainty, symptoms or unfamiliar treatment. Validate against the patient's assessment.	Short-term: Before any clinically approved transition: Identifies one immediate concern and participates in an agreed next step. Longer-term: Before the clinically approved transition or agreed follow-up: Describes helpful support and coping strategies before transition of care.	1. Listen without judgment and acknowledge specific concerns. Rationale: Feeling heard supports trust and therapeutic engagement. 2. Explain one step at a time and use teach-back with permission. Rationale: Manageable information reduces uncertainty and checks understanding. 3. Offer culturally appropriate support and refer when distress persists or safety concerns arise. Rationale: Additional support may be needed beyond bedside reassurance.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask what the learner already knows and wants to	Learning needs about the condition and care plan. Possible	Short-term: Before any clinically approved transition: Explains the	1. Assess language, literacy, hearing, vision and preferred learning method. Rationale: Accessible teaching improves	Record the baseline and reassessed

Nurses Content For educational purposes only. 5 / 8

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
understand. Assess: Use teach-back or demonstration to identify specific gaps.	factors: new information, complex treatment or limited prior teaching. Validate against the patient's assessment.	most important immediate action in their own words. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates the agreed care skills and names follow-up and warning signs.	understanding. 2. Teach a few condition-specific points using plain language and demonstration. Rationale: Focused information is easier to retain and apply. 3. Use teach-back, correct misunderstandings and provide an accessible written plan. Rationale: Checking understanding supports safer self-management. 4. Teach insulin delivery, monitoring, school support, hypoglycemia and sick-day action with age-appropriate independence. Rationale: Condition-specific teaching connects the treatment plan to safe continuing care.	finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 6 / 8

Nurses Content
Downloaded from nursescontent.com

Health education and continuing care

Teach insulin delivery, monitoring, school support, hypoglycemia and sick-day action with age-appropriate independence.

Transition only after clinical readiness is confirmed. Reconcile medicines, check access and caregiver support, and document who will review the patient and when.

Urgent warning signs

Vomiting with ketones, deep breathing, severe low glucose or altered consciousness needs emergency help.

Evaluation and conclusion

For each goal, document met, partly met or not met using observed evidence and the review time. Reassess and revise if progress is inadequate. Individual assessment, prioritized care and coordinated review guide this topic.

Quick Revision

- Autoimmune loss of insulin-producing cells creates a need for insulin replacement.

- Priority concerns: Blood glucose instability requiring close monitoring; Insufficient fluid replacement for current losses; Emotional distress affecting participation in care; Learning needs about the condition and care plan.

- Vomiting with ketones, deep breathing, severe low glucose or altered consciousness needs emergency help.

- Teach insulin delivery, monitoring, school support, hypoglycemia and sick-day action with age-appropriate independence.

Exam / viva: Which unresolved problem would delay transition, and what information must reach the next care team?

References

Nursing Fundamentals, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2024
https://wtcs.pressbooks.pub/nursingfundamentals/

Health conditions A to Z: external further reading - NHS
https://www.nhs.uk/conditions/

Nursing Pharmacology, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2023
https://wtcs.pressbooks.pub/pharmacology2e/

B.Sc. Nursing curriculum and regulatory notifications - Indian Nursing Council
https://www.indiannursingcouncil.org/gazette-notification

Original cases and care-plan wording by Nurses Content. Linked sources support further study; source terms and local clinical requirements apply. Prepared 23 November 2026. Not independently clinically peer reviewed.

Nurses Content For educational purposes only. 7 / 8

Nurses Content
Downloaded from nursescontent.com

Rights and reuse

Delivered by Nurses Content. Copyright claims apply only to original material owned by its authors. Medical facts, statutory exceptions and third-party/open licence rights remain unaffected. See the website's /policies/reuse/ page for terms.

Nurses Content For educational purposes only. 8 / 8