Nurses Content
Downloaded from nursescontent.com

Assisted personal hygiene Response monitoring

Nursing Foundation | Response monitoring

For educational purposes only.

Fictional educational example. Individualize using verified patient findings, authorized treatment and current local protocols. No real patient record or observed outcome is claimed.

Clinical assessment

Subjective: Wants to wash independently but tires quickly.

Objective findings and required checks: Needs partial help with bathing and dressing; skin is intact but folds need drying.

Physical examination and investigations: Assess task-specific ability, skin, fatigue and personal preferences.

Supply actual values, units, timing and trends during supervised application. Do not invent missing clinical data.

Clinical focus and pathophysiology

Assisted personal hygiene: Reduced stamina temporarily limits daily activities and can increase moisture-related skin problems.

Medical / surgical management

Offer a partial assisted wash, maintain privacy, check water temperature and encourage safe participation.

Illustrative drug and treatment study

Nurses Content For educational purposes only. 1 / 7

Nurses Content
Downloaded from nursescontent.com

Medicine / treatment	Purpose	Nursing checks
Emollient	Support dry skin when appropriate	Check allergy, suitability and slipping risk
barrier preparation	Protect moisture-exposed skin	Review skin response.

Treatment examples are not a prescription. Verify the authorized product, dose, route, timing, contraindications and current local product information.

Nurses Content For educational purposes only. 2 / 7

Nurses Content
Downloaded from nursescontent.com

Nursing care plan

Nursing diagnoses use original clinical problem wording. They are not the licensed NANDA-I classification. Verify any formal label or code against your institution's licensed current edition; select actual or risk diagnoses only when supported by assessment.

Context task: Compare the current assessment with the documented baseline and identify benefit, harm or lack of response.

Set individualized measurable targets and an exact review time. Select actual or risk statements only when supported by assessment.

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask which activities the person can do and where help is wanted. Assess: Assess personal-care tasks, cognition, mobility and available support.	Daily self-care needs exceed current ability. Possible factors: weakness, symptoms or reduced confidence. Validate against the patient's assessment.	Short-term: At the next scheduled reassessment, with its interval chosen for acuity: Completes a chosen self-care activity with the agreed level of assistance. Longer-term: Before the clinically approved transition or agreed follow-up: Patient or caregiver demonstrates essential care skills and knows support options.	1. Assess ability task by task and respect preferences and privacy. Rationale: Specific assessment prevents unnecessary dependence. 2. Provide only the assistance required and use suitable aids. Rationale: Supported participation protects dignity and builds independence. 3. Teach caregivers with consent and confirm realistic support at home. Rationale: Continuity depends on available skills and resources. 4. Offer a partial assisted wash, maintain privacy, check water temperature and encourage safe participation. Rationale: Reduced stamina temporarily limits daily activities and can increase moisture-related skin problems. The condition-specific plan addresses the underlying problem while response is monitored.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 3 / 7

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask about discomfort, usual mobility and ability to reposition. Assess: Inspect skin, pressure points, wound characteristics and moisture exposure.	Skin damage or vulnerability to pressure-related injury. Possible factors: pressure, moisture, impaired sensation or tissue injury. Validate against the patient's assessment.	Short-term: At the next scheduled reassessment, with its interval chosen for acuity: Skin is protected and existing damage shows no avoidable deterioration. Longer-term: Before the clinically approved transition or agreed follow-up: Patient or caregiver demonstrates the individualized skin-protection plan.	1. Assess skin and pressure risk using the local tool and clinical judgment. Rationale: Risk identification guides targeted prevention. 2. Individualize repositioning, heel offloading and pressure redistribution. Rationale: Reducing sustained pressure and shear protects tissue perfusion. 3. Use appropriate wound care and moisture management; refer for nutrition or specialist advice when needed. Rationale: Healing requires local protection and support for underlying contributors.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask about usual mobility, fatigue and help needed with activities. Assess: Assess strength, range, gait, balance and ordered movement restrictions.	Movement limited by weakness, discomfort or restriction. Possible factors: reduced strength, pain or prescribed movement limits. Validate against the patient's assessment.	Short-term: At the next scheduled reassessment, with its interval chosen for acuity: Completes a jointly agreed movement task safely with appropriate assistance. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates the home exercise or mobility plan before discharge.	1. Assess assistance needs and confirm prescribed restrictions before mobilization. Rationale: Matching support to ability prevents injury. 2. Plan graded movement, position changes and rest with physiotherapy where indicated. Rationale: Progressive activity supports function while limiting exhaustion. 3. Monitor tolerance and stop for new concerning symptoms. Rationale: Symptoms may indicate unsafe exertion or deterioration.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask what the learner already knows and wants to understand. Assess:	Learning needs about the condition and care plan. Possible factors: new	Short-term: At the next scheduled reassessment, with its interval chosen for acuity: Explains the	1. Assess language, literacy, hearing, vision and preferred learning method. Rationale: Accessible teaching improves understanding. 2. Teach a few	Record the baseline and reassessed finding, the time

Nurses Content For educational purposes only. 4 / 7

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Use teach-back or demonstration to identify specific gaps.	information, complex treatment or limited prior teaching. Validate against the patient's assessment.	most important immediate action in their own words. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates the agreed care skills and names follow-up and warning signs.	condition-specific points using plain language and demonstration. Rationale: Focused information is easier to retain and apply. 3. Use teach-back, correct misunderstandings and provide an accessible written plan. Rationale: Checking understanding supports safer self-management. 4. Teach safe hygiene aids, oral care, skin-fold drying and gradual independence. Rationale: Condition-specific teaching connects the treatment plan to safe continuing care.	and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 5 / 7

Nurses Content
Downloaded from nursescontent.com

Health education and continuing care

Teach safe hygiene aids, oral care, skin-fold drying and gradual independence.

Transition only after clinical readiness is confirmed. Reconcile medicines, check access and caregiver support, and document who will review the patient and when.

Urgent warning signs

New pressure damage, painful rash, fever or sudden functional decline needs review.

Evaluation and conclusion

For each goal, document met, partly met or not met using observed evidence and the review time. Reassess and revise if progress is inadequate. Individual assessment, prioritized care and coordinated review guide this topic.

Quick Revision

- Reduced stamina temporarily limits daily activities and can increase moisture-related skin problems.

- Priority concerns: Daily self-care needs exceed current ability; Skin damage or vulnerability to pressure-related injury; Movement limited by weakness, discomfort or restriction; Learning needs about the condition and care plan.

- New pressure damage, painful rash, fever or sudden functional decline needs review.

- Teach safe hygiene aids, oral care, skin-fold drying and gradual independence.

Exam / viva: Which observations demonstrate response, and which findings would require revising or stopping the current intervention?

References

Nursing Fundamentals, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2024
https://wtcs.pressbooks.pub/nursingfundamentals/

Nursing Pharmacology, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2023

Nurses Content For educational purposes only. 6 / 7

Nurses Content
Downloaded from nursescontent.com

https://wtcs.pressbooks.pub/pharmacology2e/

B.Sc. Nursing curriculum and regulatory notifications - Indian Nursing Council

https://www.indiannursingcouncil.org/gazette-notification

Original cases and care-plan wording by Nurses Content. Linked sources support further study; source terms and local clinical requirements apply. Prepared 23 November 2026. Not independently clinically peer reviewed.

Rights and reuse

Delivered by Nurses Content. Copyright claims apply only to original material owned by its authors. Medical facts, statutory exceptions and third-party/open licence rights remain unaffected. See the website's /policies/reuse/ page for terms.

Nurses Content For educational purposes only. 7 / 7