Nurses Content
Downloaded from nursescontent.com

Eating-disorder medical support Response monitoring

Mental Health Nursing | Response monitoring

For educational purposes only.

Fictional educational example. Individualize using verified patient findings, authorized treatment and current local protocols. No real patient record or observed outcome is claimed.

Clinical assessment

Subjective: Fear of weight gain, fatigue and anxiety around meals.

Objective findings and required checks: Medical stability, hydration and nutritional risk require specialist assessment.

Physical examination and investigations: Review observations, weight trajectory, ECG, electrolytes, glucose and individualized psychiatric assessment.

Supply actual values, units, timing and trends during supervised application. Do not invent missing clinical data.

Clinical focus and pathophysiology

Eating-disorder medical support: Restriction and malnutrition can affect cardiac, endocrine and metabolic function; refeeding needs supervision.

Medical / surgical management

Use a specialist nutrition and monitoring plan, compassionate boundaries and psychological support.

Illustrative drug and treatment study

Nurses Content For educational purposes only. 1 / 7

Nurses Content
Downloaded from nursescontent.com

Medicine / treatment	Purpose	Nursing checks
Prescribed electrolyte or vitamin replacement	Correct defined deficits within the refeeding plan	Monitor laboratories and cardiac status
prescribed psychiatric medicine	Treat selected associated symptoms	Review nutritional and cardiac safety.

Treatment examples are not a prescription. Verify the authorized product, dose, route, timing, contraindications and current local product information.

Nurses Content For educational purposes only. 2 / 7

Nurses Content
Downloaded from nursescontent.com

Nursing care plan

Nursing diagnoses use original clinical problem wording. They are not the licensed NANDA-I classification. Verify any formal label or code against your institution's licensed current edition; select actual or risk diagnoses only when supported by assessment.

Context task: Compare the current assessment with the documented baseline and identify benefit, harm or lack of response.

Set individualized measurable targets and an exact review time. Select actual or risk statements only when supported by assessment.

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask about usual diet, recent intake, weight change and food access. Assess: Assess growth or weight trends, intake record and relevant nutritional findings.	Nutritional intake below current needs. Possible factors: poor appetite, increased requirements or feeding difficulty. Validate against the patient's assessment.	Short-term: At the next scheduled reassessment, with its interval chosen for acuity: Tolerates the agreed nutrition plan without new complications. Longer-term: Before the clinically approved transition or agreed follow-up: Explains the individualized meal or feeding plan and follow-up arrangements.	1. Assess nutritional risk and confirm swallowing safety and any fasting order. Rationale: The appropriate feeding route depends on clinical safety. 2. Offer suitable small meals or prescribed feeds with dietitian input when indicated. Rationale: Individualized support improves intake and nutrient adequacy. 3. Monitor tolerance, weight or growth and metabolic findings where required. Rationale: Progress and complications guide adjustment of support. 4. Use a specialist nutrition and monitoring plan, compassionate boundaries and psychological support. Rationale: Restriction and malnutrition can affect cardiac, endocrine and metabolic function; refeeding needs supervision. The condition-specific plan addresses the underlying problem while response is monitored.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 3 / 7

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask about dizziness, chest discomfort, fatigue and cold extremities. Assess: Assess consciousness, pulse, blood pressure, capillary refill and urine trend.	Reduced circulation to vulnerable tissues. Possible factors: altered blood flow or cardiac function. Validate against the patient's assessment.	Short-term: At the next scheduled reassessment, with its interval chosen for acuity: Circulation and consciousness remain within the clinician-agreed targets. Longer-term: Before the clinically approved transition or agreed follow-up: Describes symptoms that require immediate review and the agreed follow-up plan.	1. Trend perfusion findings and escalate new chest pain, shock or altered consciousness immediately. Rationale: Poor circulation can rapidly threaten organ function. 2. Balance activity and rest and maintain the ordered monitoring schedule. Rationale: Reducing demand supports recovery while trends guide treatment. 3. Administer prescribed therapy and monitor effects and adverse reactions. Rationale: Response monitoring supports safe correction of the underlying problem.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask what worries the patient most and how they prefer to receive information. Assess: Observe distress, attention and ability to engage without making assumptions.	Emotional distress affecting participation in care. Possible factors: uncertainty, symptoms or unfamiliar treatment. Validate against the patient's assessment.	Short-term: At the next scheduled reassessment, with its interval chosen for acuity: Identifies one immediate concern and participates in an agreed next step. Longer-term: Before the clinically approved transition or agreed follow-up: Describes helpful support and coping strategies before transition of care.	1. Listen without judgment and acknowledge specific concerns. Rationale: Feeling heard supports trust and therapeutic engagement. 2. Explain one step at a time and use teach-back with permission. Rationale: Manageable information reduces uncertainty and checks understanding. 3. Offer culturally appropriate support and refer when distress persists or safety concerns arise. Rationale: Additional support may be needed beyond bedside reassurance.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask what the learner already knows and wants to understand. Assess:	Learning needs about the condition and care plan. Possible factors: new	Short-term: At the next scheduled reassessment, with its interval chosen for acuity: Explains the	1. Assess language, literacy, hearing, vision and preferred learning method. Rationale: Accessible teaching improves understanding. 2. Teach a few	Record the baseline and reassessed finding, the time

Nurses Content For educational purposes only. 4 / 7

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Use teach-back or demonstration to identify specific gaps.	information, complex treatment or limited prior teaching. Validate against the patient's assessment.	most important immediate action in their own words. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates the agreed care skills and names follow-up and warning signs.	condition-specific points using plain language and demonstration. Rationale: Focused information is easier to retain and apply. 3. Use teach-back, correct misunderstandings and provide an accessible written plan. Rationale: Checking understanding supports safer self-management. 4. Explain supervised nutrition, follow-up and support without weight stigma or unsupervised dietary targets. Rationale: Condition-specific teaching connects the treatment plan to safe continuing care.	and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 5 / 7

Nurses Content
Downloaded from nursescontent.com

Health education and continuing care

Explain supervised nutrition, follow-up and support without weight stigma or unsupervised dietary targets.

Transition only after clinical readiness is confirmed. Reconcile medicines, check access and caregiver support, and document who will review the patient and when.

Urgent warning signs

Syncope, arrhythmia, severe electrolyte disturbance, rapid deterioration or self-harm risk requires urgent care.

Evaluation and conclusion

For each goal, document met, partly met or not met using observed evidence and the review time. Reassess and revise if progress is inadequate. Individual assessment, prioritized care and coordinated review guide this topic.

Quick Revision

- Restriction and malnutrition can affect cardiac, endocrine and metabolic function; refeeding needs supervision.

- Priority concerns: Nutritional intake below current needs; Reduced circulation to vulnerable tissues; Emotional distress affecting participation in care; Learning needs about the condition and care plan.

- Syncope, arrhythmia, severe electrolyte disturbance, rapid deterioration or self-harm risk requires urgent care.

- Explain supervised nutrition, follow-up and support without weight stigma or unsupervised dietary targets.

Exam / viva: Which observations demonstrate response, and which findings would require revising or stopping the current intervention?

References

Nursing Fundamentals, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2024
https://wtcs.pressbooks.pub/nursingfundamentals/

Nursing Mental Health and Community Concepts, 2e - Open RN / WisTech Open, 2025

Nurses Content For educational purposes only. 6 / 7

Nurses Content
Downloaded from nursescontent.com

https://wtcs.pressbooks.pub/nursingmhcc/

Nursing Pharmacology, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2023

https://wtcs.pressbooks.pub/pharmacology2e/

B.Sc. Nursing curriculum and regulatory notifications - Indian Nursing Council

https://www.indiannursingcouncil.org/gazette-notification

Original cases and care-plan wording by Nurses Content. Linked sources support further study; source terms and local clinical requirements apply. Prepared 23 November 2026. Not independently clinically peer reviewed.

Rights and reuse

Delivered by Nurses Content. Copyright claims apply only to original material owned by its authors. Medical facts, statutory exceptions and third-party/open licence rights remain unaffected. See the website's /policies/reuse/ page for terms.

Nurses Content For educational purposes only. 7 / 7