Nurses Content
Downloaded from nursescontent.com

Normal labor supportive care Initial assessment

Midwifery & O.G. Nursing | Initial assessment

For educational purposes only.

Fictional educational example. Individualize using verified patient findings, authorized treatment and current local protocols. No real patient record or observed outcome is claimed.

Clinical assessment

Subjective: Painful regular contractions and a wish for continuous support.

Objective findings and required checks: Maternal/fetal observations and labor progress are assessed using the local pathway.

Physical examination and investigations: Review maternal observations, fetal monitoring, progress, hydration and preferences.

Supply actual values, units, timing and trends during supervised application. Do not invent missing clinical data.

Clinical focus and pathophysiology

Normal labor supportive care: Coordinated uterine contractions and cervical change support birth; needs vary across labor.

Medical / surgical management

Provide respectful support, comfort and mobility choices when safe, and prescribed analgesia with ongoing assessment.

Illustrative drug and treatment study

Nurses Content For educational purposes only. 1 / 7

Nurses Content
Downloaded from nursescontent.com

Medicine / treatment	Purpose	Nursing checks
Prescribed labor analgesia	Relieve pain according to informed preference	Monitor maternal/fetal effects and route-specific risks
oxytocin only when indicated	Manage a defined labor indication	Use the authorized infusion and fetal/uterine monitoring protocol.

Treatment examples are not a prescription. Verify the authorized product, dose, route, timing, contraindications and current local product information.

Nurses Content For educational purposes only. 2 / 7

Nurses Content
Downloaded from nursescontent.com

Nursing care plan

Nursing diagnoses use original clinical problem wording. They are not the licensed NANDA-I classification. Verify any formal label or code against your institution's licensed current edition; select actual or risk diagnoses only when supported by assessment.

Context task: Establish the immediate clinical priorities and verify the baseline before assuming a diagnosis.

Set individualized measurable targets and an exact review time. Select actual or risk statements only when supported by assessment.

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask about pain site, onset, severity, quality and the patient's acceptable pain level. Assess: Observe guarding, expression and effect on movement; use an appropriate pain scale.	Pain interfering with rest or activity. Possible factors: tissue irritation, injury or the underlying condition. Validate against the patient's assessment.	Short-term: After the first indicated intervention, at the documented review time: Pain decreases toward the patient-agreed target after reassessment. Longer-term: Before the clinically approved transition or agreed follow-up: Uses agreed comfort strategies and explains when new or worsening pain needs review.	1. Assess pain and associated red flags before and after treatment. Rationale: Changing pain may signal a complication and helps evaluate relief. 2. Provide comfortable positioning, reassurance and appropriate non-drug measures. Rationale: Comfort measures can reduce distress and complement prescribed treatment. 3. Give prescribed analgesia and monitor sedation, breathing and other adverse effects. Rationale: Relief should improve function without creating medication-related harm. 4. Provide respectful support, comfort and mobility choices when safe, and prescribed analgesia with ongoing assessment. Rationale: Coordinated uterine contractions and cervical change support birth; needs vary across labor. The condition-specific plan addresses the underlying problem while response is monitored.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 3 / 7

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask what worries the patient most and how they prefer to receive information. Assess: Observe distress, attention and ability to engage without making assumptions.	Emotional distress affecting participation in care. Possible factors: uncertainty, symptoms or unfamiliar treatment. Validate against the patient's assessment.	Short-term: After the first indicated intervention, at the documented review time: Identifies one immediate concern and participates in an agreed next step. Longer-term: Before the clinically approved transition or agreed follow-up: Describes helpful support and coping strategies before transition of care.	1. Listen without judgment and acknowledge specific concerns. Rationale: Feeling heard supports trust and therapeutic engagement. 2. Explain one step at a time and use teach-back with permission. Rationale: Manageable information reduces uncertainty and checks understanding. 3. Offer culturally appropriate support and refer when distress persists or safety concerns arise. Rationale: Additional support may be needed beyond bedside reassurance.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask about thirst, intake, vomiting, stools and dizziness. Assess: Monitor weight trend, intake/output, mucosa, circulation and relevant electrolytes.	Insufficient fluid replacement for current losses. Possible factors: intake below requirements or increased fluid loss. Validate against the patient's assessment.	Short-term: After the first indicated intervention, at the documented review time: Hydration and circulation improve within the agreed reassessment interval. Longer-term: Before the clinically approved transition or agreed follow-up: Patient or caregiver explains the individualized fluid plan and warning signs.	1. Measure intake and losses using age-appropriate methods. Rationale: Accurate trends guide replacement and help prevent overcorrection. 2. Offer oral fluids only when swallowing, consciousness and the treatment plan permit; otherwise implement prescribed replacement. Rationale: The safest route depends on aspiration risk and clinical severity. 3. Reassess perfusion, weight and laboratory trends after prescribed replacement. Rationale: Both continued dehydration and fluid overload require prompt recognition.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask what the learner already knows and wants to	Learning needs about the condition and care plan. Possible	Short-term: After the first indicated intervention, at the	1. Assess language, literacy, hearing, vision and preferred learning method. Rationale: Accessible teaching improves	Record the baseline and reassessed

Nurses Content For educational purposes only. 4 / 7

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
understand. Assess: Use teach-back or demonstration to identify specific gaps.	factors: new information, complex treatment or limited prior teaching. Validate against the patient's assessment.	documented review time: Explains the most important immediate action in their own words. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates the agreed care skills and names follow-up and warning signs.	understanding. 2. Teach a few condition-specific points using plain language and demonstration. Rationale: Focused information is easier to retain and apply. 3. Use teach-back, correct misunderstandings and provide an accessible written plan. Rationale: Checking understanding supports safer self-management. 4. Explain choices, breathing/comfort strategies, consent and the next stage of care. Rationale: Condition-specific teaching connects the treatment plan to safe continuing care.	finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 5 / 7

Nurses Content
Downloaded from nursescontent.com

Health education and continuing care

Explain choices, breathing/comfort strategies, consent and the next stage of care.

Transition only after clinical readiness is confirmed. Reconcile medicines, check access and caregiver support, and document who will review the patient and when.

Urgent warning signs

Abnormal fetal findings, excessive bleeding, maternal collapse, severe atypical pain or concerning labor progress needs urgent review.

Evaluation and conclusion

For each goal, document met, partly met or not met using observed evidence and the review time. Reassess and revise if progress is inadequate. Individual assessment, prioritized care and coordinated review guide this topic.

Quick Revision

- Coordinated uterine contractions and cervical change support birth; needs vary across labor.

- Priority concerns: Pain interfering with rest or activity; Emotional distress affecting participation in care; Insufficient fluid replacement for current losses; Learning needs about the condition and care plan.

- Abnormal fetal findings, excessive bleeding, maternal collapse, severe atypical pain or concerning labor progress needs urgent review.

- Explain choices, breathing/comfort strategies, consent and the next stage of care.

Exam / viva: Which finding changes the order of priorities, and what must be escalated immediately?

References

Nursing Fundamentals, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2024
https://wtcs.pressbooks.pub/nursingfundamentals/

Nurses Content For educational purposes only. 6 / 7

Nurses Content
Downloaded from nursescontent.com

Maternal-Newborn Nursing: external further reading - OpenStax / Rice University
https://openstax.org/books/maternal-newborn-nursing/pages/preface

Nursing Pharmacology, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2023
https://wtcs.pressbooks.pub/pharmacology2e/

B.Sc. Nursing curriculum and regulatory notifications - Indian Nursing Council
https://www.indiannursingcouncil.org/gazette-notification

Original cases and care-plan wording by Nurses Content. Linked sources support further study; source terms and local clinical requirements apply. Prepared 23 November 2026. Not independently clinically peer reviewed.

Rights and reuse

Delivered by Nurses Content. Copyright claims apply only to original material owned by its authors. Medical facts, statutory exceptions and third-party/open licence rights remain unaffected. See the website's /policies/reuse/ page for terms.

Nurses Content For educational purposes only. 7 / 7