Nurses Content
Downloaded from nursescontent.com

Postpartum hemorrhage Patient & caregiver education

Midwifery & O.G. Nursing | Patient & caregiver education

For educational purposes only.

Fictional educational example. Individualize using verified patient findings, authorized treatment and current local protocols. No real patient record or observed outcome is claimed.

Clinical assessment

Subjective: Feels faint and weak; bleeding is heavier than expected.

Objective findings and required checks: Blood loss and circulatory changes require an immediate obstetric emergency response.

Physical examination and investigations: Quantify loss, assess uterine tone and circulation, and obtain ordered blood count/coagulation and compatibility tests.

Supply actual values, units, timing and trends during supervised application. Do not invent missing clinical data.

Clinical focus and pathophysiology

Postpartum hemorrhage: Uterine atony, retained tissue, trauma or coagulation problems can cause serious postpartum blood loss.

Medical / surgical management

Activate the hemorrhage protocol, assess causes and perform authorized emergency measures while preparing prescribed treatment.

Illustrative drug and treatment study

Nurses Content For educational purposes only. 1 / 7

Nurses Content
Downloaded from nursescontent.com

Medicine / treatment	Purpose	Nursing checks
Prescribed uterotonic	Treat uterine atony when indicated	Check contraindications and response
prescribed tranexamic acid/blood products	Support indicated hemorrhage treatment	Verify timing, indication and reaction monitoring under protocol.

Treatment examples are not a prescription. Verify the authorized product, dose, route, timing, contraindications and current local product information.

Nurses Content For educational purposes only. 2 / 7

Nurses Content
Downloaded from nursescontent.com

Nursing care plan

Nursing diagnoses use original clinical problem wording. They are not the licensed NANDA-I classification. Verify any formal label or code against your institution's licensed current edition; select actual or risk diagnoses only when supported by assessment.

Context task: Assess understanding, demonstrate a condition-specific skill and verify learning using teach-back.

Set individualized measurable targets and an exact review time. Select actual or risk statements only when supported by assessment.

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask about bleeding, dizziness, bruising and relevant medicines. Assess: Assess blood loss, circulation, consciousness and ordered coagulation or blood counts.	Bleeding or vulnerability to further blood loss. Possible factors: tissue injury, altered coagulation or treatment effects. Validate against the patient's assessment.	Short-term: By the end of the planned teaching session: Blood loss is recognized promptly and circulation is supported by the emergency or treatment plan. Longer-term: Before the clinically approved transition or agreed follow-up: Identifies bleeding warning signs and medicine-related precautions.	1. Quantify visible loss where possible and escalate active significant bleeding immediately. Rationale: Underestimating blood loss can delay life-saving care. 2. Prepare ordered access, investigations and treatment while monitoring perfusion. Rationale: Coordinated response supports rapid stabilization. 3. Avoid unnecessary traumatic procedures and apply appropriate bleeding precautions. Rationale: Reducing additional injury limits preventable blood loss. 4. Activate the hemorrhage protocol, assess causes and perform authorized emergency measures while preparing prescribed treatment. Rationale: Uterine atony, retained tissue, trauma or coagulation problems can cause serious postpartum blood loss. The condition-specific plan addresses the underlying problem while response is monitored.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 3 / 7

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
Ask: Ask about thirst, intake, vomiting, stools and dizziness. Assess: Monitor weight trend, intake/output, mucosa, circulation and relevant electrolytes.	Insufficient fluid replacement for current losses. Possible factors: intake below requirements or increased fluid loss. Validate against the patient's assessment.	Short-term: By the end of the planned teaching session: Hydration and circulation improve within the agreed reassessment interval. Longer-term: Before the clinically approved transition or agreed follow-up: Patient or caregiver explains the individualized fluid plan and warning signs.	1. Measure intake and losses using age-appropriate methods. Rationale: Accurate trends guide replacement and help prevent overcorrection. 2. Offer oral fluids only when swallowing, consciousness and the treatment plan permit; otherwise implement prescribed replacement. Rationale: The safest route depends on aspiration risk and clinical severity. 3. Reassess perfusion, weight and laboratory trends after prescribed replacement. Rationale: Both continued dehydration and fluid overload require prompt recognition.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask about dizziness, chest discomfort, fatigue and cold extremities. Assess: Assess consciousness, pulse, blood pressure, capillary refill and urine trend.	Reduced circulation to vulnerable tissues. Possible factors: altered blood flow or cardiac function. Validate against the patient's assessment.	Short-term: By the end of the planned teaching session: Circulation and consciousness remain within the clinician-agreed targets. Longer-term: Before the clinically approved transition or agreed follow-up: Describes symptoms that require immediate review and the agreed follow-up plan.	1. Trend perfusion findings and escalate new chest pain, shock or altered consciousness immediately. Rationale: Poor circulation can rapidly threaten organ function. 2. Balance activity and rest and maintain the ordered monitoring schedule. Rationale: Reducing demand supports recovery while trends guide treatment. 3. Administer prescribed therapy and monitor effects and adverse reactions. Rationale: Response monitoring supports safe correction of the underlying problem.	Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.
Ask: Ask what the learner already knows and wants to	Learning needs about the condition and care plan. Possible	Short-term: By the end of the planned teaching session: Explains the	1. Assess language, literacy, hearing, vision and preferred learning method. Rationale: Accessible teaching improves	Record the baseline and reassessed

Nurses Content For educational purposes only. 4 / 7

Nurses Content
Downloaded from nursescontent.com

Assessment: subjective / objective	Nursing diagnosis: original problem wording	Short- and longer-term goals	Nursing interventions with rationale	Evaluation
understand. Assess: Use teach-back or demonstration to identify specific gaps.	factors: new information, complex treatment or limited prior teaching. Validate against the patient's assessment.	most important immediate action in their own words. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates the agreed care skills and names follow-up and warning signs.	understanding. 2. Teach a few condition-specific points using plain language and demonstration. Rationale: Focused information is easier to retain and apply. 3. Use teach-back, correct misunderstandings and provide an accessible written plan. Rationale: Checking understanding supports safer self-management. 4. After stabilization explain anemia/recovery care, recurrent bleeding warning signs and follow-up. Rationale: Condition-specific teaching connects the treatment plan to safe continuing care.	finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement.

Nurses Content For educational purposes only. 5 / 7

Nurses Content
Downloaded from nursescontent.com

Health education and continuing care

After stabilization explain anemia/recovery care, recurrent bleeding warning signs and follow-up.

Transition only after clinical readiness is confirmed. Reconcile medicines, check access and caregiver support, and document who will review the patient and when.

Urgent warning signs

Ongoing heavy loss, shock, collapse or reduced consciousness requires immediate senior emergency care.

Evaluation and conclusion

For each goal, document met, partly met or not met using observed evidence and the review time. Reassess and revise if progress is inadequate. Individual assessment, prioritized care and coordinated review guide this topic.

Quick Revision

- Uterine atony, retained tissue, trauma or coagulation problems can cause serious postpartum blood loss.

- Priority concerns: Bleeding or vulnerability to further blood loss; Insufficient fluid replacement for current losses; Reduced circulation to vulnerable tissues; Learning needs about the condition and care plan.

- Ongoing heavy loss, shock, collapse or reduced consciousness requires immediate senior emergency care.

- After stabilization explain anemia/recovery care, recurrent bleeding warning signs and follow-up.

Exam / viva: What should the learner demonstrate or explain before you consider the teaching effective?

References

Nursing Fundamentals, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2024
https://wtcs.pressbooks.pub/nursingfundamentals/

Maternal-Newborn Nursing: external further reading - OpenStax / Rice University
https://openstax.org/books/maternal-newborn-nursing/pages/preface

Nurses Content For educational purposes only. 6 / 7

Nurses Content
Downloaded from nursescontent.com

Nursing Pharmacology, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2023
https://wtcs.pressbooks.pub/pharmacology2e/

B.Sc. Nursing curriculum and regulatory notifications - Indian Nursing Council
https://www.indiannursingcouncil.org/gazette-notification

Original cases and care-plan wording by Nurses Content. Linked sources support further study; source terms and local clinical requirements apply. Prepared 23 November 2026. Not independently clinically peer reviewed.

Rights and reuse

Delivered by Nurses Content. Copyright claims apply only to original material owned by its authors. Medical facts, statutory exceptions and third-party/open licence rights remain unaffected. See the website's /policies/reuse/ page for terms.

Nurses Content For educational purposes only. 7 / 7