Nurses Content
Downloaded from nursescontent.com



# Sepsis recognition and initial care Transition & follow-up

Adult Health Nursing | Transition & follow-up

**For educational purposes only.**

Fictional educational example. Individualize using verified patient findings, authorized treatment and current local protocols. No real patient record or observed outcome is claimed.

## Clinical assessment

Subjective: Feels extremely unwell with weakness and reduced urine.

Objective findings and required checks: Abnormal observations and possible organ dysfunction trigger urgent clinical review.

Physical examination and investigations: Use the local sepsis assessment pathway; obtain indicated lactate, cultures, blood tests and source investigations.

Supply actual values, units, timing and trends during supervised application. Do not invent missing clinical data.

## Clinical focus and pathophysiology

Sepsis recognition and initial care: A dysregulated response to infection can impair perfusion and damage organs.

## Medical / surgical management

Escalate urgently, monitor closely and deliver prescribed antimicrobial, fluid and source-control preparations.

## Illustrative drug and treatment study



Nurses Content For educational purposes only. 1 / 7

Nurses Content
Downloaded from nursescontent.com



| Medicine / treatment | Purpose | Nursing checks |
|---|---|---|
| Prescribed antimicrobial | Treat suspected serious infection | Check allergy, obtain cultures without harmful delay and monitor response |
| prescribed IV fluid | Support assessed circulatory needs | Reassess perfusion and overload risk. |

Treatment examples are not a prescription. Verify the authorized product, dose, route, timing, contraindications and current local product information.



Nurses Content For educational purposes only. 2 / 7

Nurses Content
Downloaded from nursescontent.com



# Nursing care plan

Nursing diagnoses use original clinical problem wording. They are not the licensed NANDA-I classification. Verify any formal label or code against your institution's licensed current edition; select actual or risk diagnoses only when supported by assessment.

Context task: Check unresolved risks, medicine reconciliation, practical support and follow-up before transition.

Set individualized measurable targets and an exact review time. Select actual or risk statements only when supported by assessment.

| Assessment: subjective / objective | Nursing diagnosis: original problem wording | Short- and longer-term goals | Nursing interventions with rationale | Evaluation |
|---|---|---|---|---|
| Ask: Ask about dizziness, chest discomfort, fatigue and cold extremities. Assess: Assess consciousness, pulse, blood pressure, capillary refill and urine trend. | Reduced circulation to vulnerable tissues. Possible factors: altered blood flow or cardiac function. Validate against the patient's assessment. | Short-term: Before any clinically approved transition: Circulation and consciousness remain within the clinician-agreed targets. Longer-term: Before the clinically approved transition or agreed follow-up: Describes symptoms that require immediate review and the agreed follow-up plan. | 1. Trend perfusion findings and escalate new chest pain, shock or altered consciousness immediately. Rationale: Poor circulation can rapidly threaten organ function. 2. Balance activity and rest and maintain the ordered monitoring schedule. Rationale: Reducing demand supports recovery while trends guide treatment. 3. Administer prescribed therapy and monitor effects and adverse reactions. Rationale: Response monitoring supports safe correction of the underlying problem. 4. Escalate urgently, monitor closely and deliver prescribed antimicrobial, fluid and source-control preparations. Rationale: A dysregulated response to infection can impair perfusion and damage organs. The condition-specific plan addresses the underlying problem while response is monitored. | Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement. |



Nurses Content For educational purposes only. 3 / 7

Nurses Content
Downloaded from nursescontent.com



| Assessment: subjective / objective | Nursing diagnosis: original problem wording | Short- and longer-term goals | Nursing interventions with rationale | Evaluation |
|---|---|---|---|---|
| Ask: Ask about exposure, wound symptoms and recent invasive devices. Assess: Inspect relevant sites and monitor systemic findings and investigation trends. | Vulnerability to infection or worsening infection. Possible factors: disrupted barriers, exposure or reduced host defence. Validate against the patient's assessment. | Short-term: Before any clinically approved transition: Aseptic care is maintained and any new signs of infection are escalated promptly. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates relevant hygiene and explains the infection-prevention plan. | 1. Use hand hygiene and risk-based standard or transmission precautions. Rationale: Interrupting transmission protects the patient, staff and other patients. 2. Review invasive device need with the team and maintain indicated aseptic care. Rationale: Unnecessary devices and barrier breaks increase infection exposure. 3. Monitor for systemic deterioration and administer prescribed antimicrobials on schedule. Rationale: Early escalation and timely indicated treatment reduce avoidable harm. | Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement. |
| Ask: Ask about thirst, intake, vomiting, stools and dizziness. Assess: Monitor weight trend, intake/output, mucosa, circulation and relevant electrolytes. | Insufficient fluid replacement for current losses. Possible factors: intake below requirements or increased fluid loss. Validate against the patient's assessment. | Short-term: Before any clinically approved transition: Hydration and circulation improve within the agreed reassessment interval. Longer-term: Before the clinically approved transition or agreed follow-up: Patient or caregiver explains the individualized fluid plan and warning signs. | 1. Measure intake and losses using age-appropriate methods. Rationale: Accurate trends guide replacement and help prevent overcorrection. 2. Offer oral fluids only when swallowing, consciousness and the treatment plan permit; otherwise implement prescribed replacement. Rationale: The safest route depends on aspiration risk and clinical severity. 3. Reassess perfusion, weight and laboratory trends after prescribed replacement. Rationale: Both continued dehydration and fluid overload require prompt recognition. | Record the baseline and reassessed finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement. |
| Ask: Ask what the learner already knows and wants to | Learning needs about the condition and care plan. Possible | Short-term: Before any clinically approved transition: Explains the | 1. Assess language, literacy, hearing, vision and preferred learning method. Rationale: Accessible teaching improves | Record the baseline and reassessed |



Nurses Content For educational purposes only. 4 / 7

Nurses Content
Downloaded from nursescontent.com



| Assessment: subjective / objective | Nursing diagnosis: original problem wording | Short- and longer-term goals | Nursing interventions with rationale | Evaluation |
|---|---|---|---|---|
| understand. Assess: Use teach-back or demonstration to identify specific gaps. | factors: new information, complex treatment or limited prior teaching. Validate against the patient's assessment. | most important immediate action in their own words. Longer-term: Before the clinically approved transition or agreed follow-up: Demonstrates the agreed care skills and names follow-up and warning signs. | understanding. 2. Teach a few condition-specific points using plain language and demonstration. Rationale: Focused information is easier to retain and apply. 3. Use teach-back, correct misunderstandings and provide an accessible written plan. Rationale: Checking understanding supports safer self-management. 4. After stabilization explain treatment, recovery needs and urgent return signs. Rationale: Condition-specific teaching connects the treatment plan to safe continuing care. | finding, the time and whether this goal is met, partly met or not met. If not met, reassess causes and revise or escalate. Do not record an unobserved improvement. |



Nurses Content For educational purposes only. 5 / 7

Nurses Content
Downloaded from nursescontent.com



## Health education and continuing care

After stabilization explain treatment, recovery needs and urgent return signs.

Transition only after clinical readiness is confirmed. Reconcile medicines, check access and caregiver support, and document who will review the patient and when.

## Urgent warning signs

**Shock, altered consciousness, severe respiratory distress or rapidly falling urine output requires immediate emergency team response.**

## Evaluation and conclusion

For each goal, document met, partly met or not met using observed evidence and the review time. Reassess and revise if progress is inadequate. Individual assessment, prioritized care and coordinated review guide this topic.

## Quick Revision

- A dysregulated response to infection can impair perfusion and damage organs.

- Priority concerns: Reduced circulation to vulnerable tissues; Vulnerability to infection or worsening infection; Insufficient fluid replacement for current losses; Learning needs about the condition and care plan.

- Shock, altered consciousness, severe respiratory distress or rapidly falling urine output requires immediate emergency team response.

- After stabilization explain treatment, recovery needs and urgent return signs.

**Exam / viva: Which unresolved problem would delay transition, and what information must reach the next care team?**

## References

Nursing Fundamentals, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2024
https://wtcs.pressbooks.pub/nursingfundamentals/



Nurses Content For educational purposes only. 6 / 7

Nurses Content
Downloaded from nursescontent.com



Health Alterations - Open RN / WisTech Open, 2025
https://wtcs.pressbooks.pub/healthalts/

Nursing Pharmacology, 2e - Ernstmeyer & Christman (editors), Open RN / WisTech Open, 2023
https://wtcs.pressbooks.pub/pharmacology2e/

B.Sc. Nursing curriculum and regulatory notifications - Indian Nursing Council
https://www.indiannursingcouncil.org/gazette-notification

Original cases and care-plan wording by Nurses Content. Linked sources support further study; source terms and local clinical requirements apply. Prepared 23 November 2026. Not independently clinically peer reviewed.

## Rights and reuse

Delivered by Nurses Content. Copyright claims apply only to original material owned by its authors. Medical facts, statutory exceptions and third-party/open licence rights remain unaffected. See the website's /policies/reuse/ page for terms.



Nurses Content For educational purposes only. 7 / 7